

Review

Oxytocin-messages via the cerebrospinal fluid: Behavioral effects; a review

Jan G. Veening ^{a,b,d,*}, Trynke de Jong ^c, Henk P. Barendregt ^d

^a Department of Anatomy (109), UMC St Radboud, Nijmegen, The Netherlands

^b Department of Psychopharmacology, UIPS, Univ. of Utrecht, The Netherlands

^c Department of Biology, 3386 Spieth Hall, Univ. of California, Riverside, CA 92521, USA

^d Fac. of Science, Radboud University, Nijmegen, The Netherlands

ARTICLE INFO

Article history:

Received 12 November 2009

Received in revised form 21 April 2010

Accepted 4 May 2010

Keywords:

Cerebrospinal fluid

Oxytocin

Behavioral state

Non-synaptic communication

Dendritic release

Olfaction

ABSTRACT

The cerebrospinal fluid (CSF) usually is considered as a protective ‘nutrient and waste control’ system for the brain. Recent findings suggest, however, that the composition of CSF is actively controlled and may play an influential role in the changes in brain activity, underlying different behavioral states. In the present review, we present an overview of available data concerning the release of oxytocin into the CSF, the location of the oxytocin-receptive brain areas and the behavioral effects of intracerebroventricular oxytocin. About 80% of the oxytocin-receptive areas are located close to the ventricular or subarachnoid CSF, including the hypothalamic ‘Behavior Control Column’ (L.W.Swanson, 2003). As a conclusion we suggest that ‘CSF-oxytocin’ contributes considerably to the non-synaptic communication processes involved in hypothalamic-, brainstem- and olfactory brain areas and behavioral states and that the flowing CSF is used as a ‘broadcasting system’ to send coordinated messages to a wide variety of nearby and distant brain areas.

© 2010 Elsevier Inc. All rights reserved.

Contents

1. Introduction	194
2. Oxytocin and CSF	194
2.1. Peripheral and CSF- OT-levels are regulated differentially	194
2.2. General neuroanatomy of the oxytocin system	195
2.3. Diffusion and dendritic release from PVH	195
2.4. Supra-ependymal release	195
2.5. Synchronisation and cross-talk of OT neurons	197
2.6. OT released into the CSF: are the levels functionally relevant?	198
2.7. Oxytocin receptors and the CSF	198
3. Behavioral effects of oxytocin are elicited in the CSF	200
3.1. Social behavior	200
3.2. Sexual behavior	201
3.3. Parental behavior	201
3.4. Food intake	201
3.5. Grooming behavior	201
3.6. Pain	202
4. CSF-oxytocin may affect various parts of the CNS	202
4.1. OT- effects on hypothalamic ‘Behavioral Control Column’ (Nr 1 in Fig. 5, Fig. 6)	202
4.2. OT- effects on brainstem (‘core’- and ‘paracore’-) areas (Fig. 5, nr 2)	202
4.3. OT- effects on cortical areas (Fig. 5, nr 3)	202
4.4. OT- effects on olfactory areas (Fig. 5, nr 4)	203
4.5. OT- effects on olfactory sensory neurons (OSNs) (Fig. 5, nr 5)	203
5. Conclusions and future directions	204
References	204

* Corresponding author. Department of Anatomy, 109, UMC St Radboud, PO Box 9101, 6500 HB, Nijmegen, The Netherlands. Tel.: +31 243614298.
E-mail address: j.veening@anat.umcn.nl (J.G. Veening).

1. Introduction

Animal and human behavior can be described as 'goal-directed' with a succession of different behaviors leading to different goals (feeding and drinking, or 'social activities' like aggression or sex) over time. Such changes in behavior require adaptive physiological mechanisms and complex changes in neural activity in a wide variety of brain areas and have been described as switches of the 'behavioral state' of the organism. Such states regulate the input–output repertoire of the organism and have been described and defined as motivational, emotional or mental states [1–6] in animal and human behavioral studies. Obviously, the hypothalamus, embedded in the limbic system is necessarily involved when motivational states change [1,3,6–8], but other brain areas, ranging from cortical regions to spinal cord, must be involved as well in order to accomplish the coordinated adaptations in the neural (sensory, motor and autonomic) as well as hormonal systems, underlying goal-directed behavior.

Since the early eighties, the occurrence of 'non-synaptic communication' or 'volume transmission' has been described as a new mechanism in addition to 'wiring transmission' [9–13]. 'Volume transmission' has been proposed as a mechanism to change neuronal activity in more than a single brain area, for an extended period of time [1,7,8,14–23], which are basic requirements for motivational changes. Many descending neuropeptidergic fiber systems containing mostly unmyelinated varicose fibers and lacking synaptic contacts [1,6–8,20,22,24–26] are in an excellent position for changing the weight factors in the neural network involved in keeping or switching between behavioral states. Remarkably, many of these descending fibers are running close to the ventricular system [24–27], which raises the question in how far the cerebrospinal fluid (CSF), flowing through the ventricular system and the arachnoid space surrounding the brain, could be involved in behavioral changes.

In early studies of Cushing [28] the term 'third circulation' was coined for the CSF, flowing caudally through as well as around the CNS. Friedman and Friedman [29] showed already that oxytocin-levels in the CSF were sufficiently high to induce contractions of the rabbit uterus, which makes it legitimate to ask in how far the circulating CSF can play a role in physiological and behavioral changes and - states.

Previously, Sowards and Sowards [30,31] have argued, in our view convincingly, that vasopressin and corticotropin-releasing-factor (CRF) are released into the CSF in order to induce their behavioral effects via activation of the appropriate receptors in distant brain areas. In a previous paper [23] we have discussed the available evidence that the flow of the CSF through and around the CNS provides a fast and specific substrate for sending neuropeptidergic as well as other 'messages' to appropriate receptive brain areas. Especially the local release of melatonin into the dorsal part of the third ventricle and the release of 'gonadotropin hormone releasing hormone' (GnRH) into the basal part of the 3rd ventricle, each with their own specific target brain areas [32–37] strongly support the hypothesis that the CSF may be involved in 'broadcasting messages' as a special kind of 'volume transmission'.

In the present review we intend to explore the available evidence that oxytocin (OT)-producing neurons are using the CSF as a 'broadcasting system' to communicate with a variety of other nearby as well as distant brain areas. For the assumption that OT distributed via the CSF is relevant for behavioral changes, at least the following kinds of supporting evidence are necessary:

- OT can be released into the CSF, 'to go with the flow';
- OT-levels in the CSF are regulated independently and different from the peripheral levels in the blood;
- OT-levels in the CSF are sufficiently high to activate OT-receptive areas alongside the ventricular or arachnoid spaces;

- OT-receptive 'target'-areas are located alongside the ventricular and arachnoid spaces of the brain;
- OT-administration into the CSF induces behavioral effects not obtained by peripheral administration;
- behavioral effects of intra-cerebro-ventricular (icv) -OT-administration can be linked, anatomically and/or functionally, to the OT-receptive brain areas.

For each of these aspects, the available evidence will be presented and discussed in the next sections of the present review.

2. Oxytocin and CSF

Oxytocin is a nonapeptide, synthesized by Du Vigneaud in 1953, after its isolation and clarification [38]. Since then, OT has been implicated in a dazzling variety of widely diverging functions. Many reviews are available summarising and highlighting the relevant specific findings in detail: in each of the years 2007 and 2008 at least 20 reviews appeared focusing on specific OT-functions.

2.1. Peripheral and CSF- OT-levels are regulated differentially

OT has been detected in the cerebrospinal fluid of many species, although the levels vary considerably both diurnally and between species [39–43].

It has been shown extensively that OT levels in the CSF are different from plasma OT levels and react very differently upon natural or experimental challenges. Natural and experimentally manipulated circadian rhythms are much more visible in the CSF than in the blood plasma [39–42,44]. Effects of pregnancy and labor in humans [45], of parturition in sheep or rats [46,47], of hand-milking in the goat [48] or of suckling in the guinea pig [49] are very different in CSF compared to plasma levels of OT. Experimental challenges like opioid or naltrexone administration in rats [50] and sheep [51], partner preference test [52], stress [53], vagino-cervical stimulation in the goat [54], electrical stimulation of the hypothalamus or neural lobe in the rat [55], chemical stimulation of the paraventricular hypothalamic nucleus (PVH) by the excitatory amino acid glutamate [56] and hypophysectomy [57] all affect OT-levels in CSF differently from plasma OT-levels. Many additional findings are available from the literature to draw the conclusion that CSF and peripheral levels of OT are not only different but, more importantly, are controlled by different largely independent, mechanisms [42]. The finding that peripheral OT-levels contribute only 0.002% to the OT-CSF-levels [58], supports the functional independency. Peripheral levels may not be completely independent, however, since CSF-OT may play a regulatory role in the control of peripheral OT levels [48].

In addition to the differentially regulated levels, the half-life values of OT are completely different: about 28 min in CSF compared to 1–2 min in the blood plasma [49,58–60], which suggests that the behavioral effects of OT inside the CNS use a radically different time-scale compared to the peripheral effects on specific organs.

From the multiple functional differences in the control of CSF and plasma OT levels, we propose that it is certainly allowed to explore the possibility that ventricular release of OT into the CSF may compose a message for other parts of the brain, because, once released, it may 'follow the flow' and reach many widespread brain areas within minutes. In the following parts of this review, the possible origins and destinations of the 'CSF-OT-message' will be discussed.

In summary: CSF-OT levels are not controlled by peripheral systemic levels but by independent mechanisms controlling both levels and temporary peaks in the CSF. Combined with the about 20 times longer half-time values of OT in the CSF, this suggests that independent 'OT-messages' to other brain areas are possible, without involvement of peripheral effects on specific organs.

2.2. General neuroanatomy of the oxytocin system

The production and subsequent central or peripheral release of OT is executed by partly overlapping and interconnected groups of neurons. The majority of OT-producing neurons is located in the paraventricular hypothalamic nuclei (PVH), with a dorsal extension into the bed nucleus of the stria terminalis, and in the supraoptic hypothalamic nuclei (SON). Several additional perivascular clusters of OT-immunoreactive (OT-IR) neurons can be observed consistently in the rostral hypothalamus, 'scattered' between the SON and PVH. These clusters are interesting because of the special role of the perivascular spaces for the rapid distribution of CSF-messages [61–64].

Parvocellular neurons in the dorsal part of the PVH release OT from their axons in distant CNS sites [65,66]. Interestingly, many of these axonal projections reach areas bordering, surrounding and/or functionally related to the ventricular system [65–71]. Magnocellular neurons in the paraventricular and supraoptic nuclei of the hypothalamus release OT directly from their axons into the blood stream via the neurohypophysis. Many morphological and functional aspects of both types of OT-neurons (projections, afferent connections, co-localisation with other neuropeptides, plasticity, etc) have been discussed in a lengthy series of review papers since 1983 [66], and can be left out of focus for the present purpose.

Production of OT occurs in 2 hypothalamic cell groups (the paraventricular and the supraoptic nuclei) as well as in a few perivascular clusters in between. Both brain areas border the CSF directly, (at the third ventricle or the arachnoid space, respectively) and contain magnocellular neurons that release OT via the pituitary directly into the systemic circulation. The parvocellular OT-neurons of the PVH send their axonal projections to many distant brain areas, including the spinal cord, often at a small distance from the ventricular system.

2.3. Diffusion and dendritic release from PVH

Hoistad et al. [72] have convincingly shown that neuropeptides like beta-endorphin, released at a distance, eventually appear in the CSF, with a peak-delay of about half an hour. Their finding shows two important matters: on the one hand that axonally-released neuropeptides eventually reach the CSF through diffusion and/or extracellular fluid (ECF-) flow [73]; on the other hand that peptidase-levels in the ECF and CSF are apparently too low to prevent the appearance of peptides in the CSF, half an hour later. Apparently, in the CNS extracellular neuropeptides are distributed and may remain effective over considerable distances for an extended period of time. On the other hand, however, the appearance of neuropeptides into the CSF is slow and levels remain generally low, when released at such distances from the ventricular system. In agreement with this suggestion, a recent paper showed that only about 10 to 15% of the ECF is drained into the CSF [74]. As a mechanism for 'switching behavioral states' such ECF-CSF-exchange is probably too weak and too slow but 'sustaining effects' cannot be excluded. Therefore, the notion of Landgraf & Neumann [69] may be generally right, that 'neuropeptide levels in the CSF essentially reflect the more or less "global" activity of the corresponding neuropeptide in the brain'. However, in the case of OT, a 5-fold daily increase in primate CSF-OT [40] has been observed. Without specific release mechanisms, such strong fluctuations cannot be explained by 'axonal/terminal release at a distance'. Interestingly, over the last few years, appropriate mechanisms have been described and investigated that are capable of releasing large amounts of OT specifically into the CSF.

This fascinating role might be played by dendritic release and subsequent diffusion of oxytocin in the PVH and the SON [69,75–77]. Inside the PVH, it is remarkable to observe how the dendrites of the magnocellular (as well as other) parts of the PVH generally run in

parallel to the outer borders of the PVH(-subdivisions) and show a strong preference to extend into the central-medial parts of the PVH (Fig. 1). Some of the medially directed dendrites not only reach and penetrate (Fig. 1A,B) the ventricular lining but some of them have been observed to cross the dorsal side of the third ventricle, potentially to influence the contralateral PVH as well (Fig. 1B) [78–81]. In combination with the lateral cell-free border zone, along the ventrolateral side of the PVH, which limits diffusion in a ventrolateral direction (pers.obs.), this particular 'internally directed' dendritic organisation strongly suggests a specific function in the 'internal' communication inside the PVH as well as towards the CSF (Fig. 2). The dendrites of the OT-neurons inside the SON show an even stronger preference to extend in a ventral direction, towards the subarachnoid space. All of these magnocellular dendrites have been shown to release OT [76,82,83] and have been described as 'prototypic receiver/transmitters' [84,85] (Fig. 3).

Since OT has a strong excitatory effect on the activity of (surrounding) OT-neurons [84,86,87], dendritic exocytosis turned out to form the basis for a chain reaction of massive OT-activation and release of OT inside the PVH and SON leading to a thousand-fold increase of extracellular OT [69,75,76], inevitably followed by increases in CSF-levels because of the missing ECF-CSF-barriers. This specific type of release may contribute to coordinate the action of large groups of OT neurons via positive feedback mechanisms, and may also play a role to functionally separate the peripheral release, in the neural lobe of the pituitary, from the central effects inside the hypothalamic nuclei. Dendritic release itself occurs at a millisecond time scale, but the size of the effects on surrounding neurons is strongly depending on the hormonal state of the animal [88,89] and appears to be controlled by the presence of oestradiol and estrogen receptivity [89,90], by specific changes in neuron-glia interactions, by numbers of plasmalemma juxtapositions and by priming effects [75,76,84,91–93]. In addition, dendritic release of OT may be self-sustaining and long-lasting (up to 90 min). These dendritic release mechanisms are in a complex way supported by specific priming effects [93], and by the twentyfold elongated half-life values in CSF compared to blood plasma [49,58–60]. Mechanisms controlling the receptive properties of surrounding neurons by insertion of receptor proteins into postsynaptic membranes, [84] contribute to the complexity.

For the cellular mechanisms involved in the differential regulation of dendritic versus axonal release of OT, the reader is referred to an extensive series of recent review papers [69,76,83,85,94].

In summary: neuropeptides released elsewhere in the brain will eventually appear in the CSF, sometimes after considerable delay. The thousand-fold increases of OT-levels in the PVH and SON as a result of massive and coordinated dendritic release mechanisms will, however, strongly and almost immediately affect OT-levels in the CSF. Since the dendritic and the terminal release mechanisms of the magnocellular neurons appear to be functionally independent, dendritic release mechanisms are the best candidates to regulate CSF-OT levels independently from systemic release.

2.4. Supra-ependymal release

In addition to the dendritic release followed by diffusion of OT into the CSF, another source of OT-release is composed by the 'supra- and sub-ependymal structures'.

It has been observed that OT-IR dendrites run subependymally along the 3rd ventricular wall, in a rostral or caudal direction [95], and that supraependymal fibers run along the ventricular surface to terminate in the supraependymal plexus on the floor of the 3rd ventricle [96]. Large numbers of 'neural nitric oxide synthase (nNOS)-positive cerebrospinal fluid-contacting neurons' have been observed in the rat, with somas, dendrites and axons situated at sub- and supraependymal levels along the 3rd ventricular wall. 88% of these




Fig. 1. Shows the location of paraventricular hypothalamic neurons and dendrites with their characteristic extensions in the medial direction, towards the ventricular surface, with exceptionally few dendrites leaving the confines of the paraventricular hypothalamic nucleus. Panel A is reprinted from: Rho and Swanson [81], with permission of the Society of Neuroscience. Abbreviation: pv: periventricular part of the PVH, bordering the 3rd ventricle; B, C and D clearly show the medialward preference of the dendritic extensions towards the 3rd ventricle on the left. In B the 2 upper medial neurons show dendritic extensions that may cross the tip of the 3rd ventricle to extend to (dendrites from) the contralateral paraventricular nucleus, maybe contributing to bilateral PVH-synchronisation. B,C and D are reprinted from: Armstrong et al. [80], with permission of Elsevier.

neurons were OT-IR [97]. In addition, in mice OT-immunoreactive cells with cilia extending into the 3rd ventricle from the lower part of the anterior ventricular wall near the suprachiasmatic nucleus, showed a significant daily rhythm in immunostaining, regulated by environmental lighting [44]. These effects were 'OT-specific' since no effects on calbindin or vasopressin immunostaining were observed.

Combining these data with older data concerning neurophysin (a carrier protein involved in the axonal transport of PVH-OT) [98], with the supposed general role of CSF-contacting neurons as reviewed by

[99,100], and with some of the data concerning the magnocellular PVH-neurons, as discussed above, it is only a small step to suggest that oxytocinergic fibers and 'CSF-contacting-neurons' surrounding the ependymal layer of the 3rd ventricle, may directly and actively contribute to the OT-levels in the CSF.

There are no data available to estimate the size, or importance, of this contribution relative to the amounts released by the magnocellular PVH- and SON dendrites. It may be questioned, however, whether it makes sense to discuss these contributions as separate entities. The




Fig. 2. Figure is reprinted from Dubois-Dauphin et al. [163], with permission of Elsevier. A and B show clearly how close OT-immunoreactive neurons are bordering the third ventricle in the guinea pig, in an excellent position to release their dendritic contents into the CSF. In addition, B shows many OT-IR fibers penetrating the ependymal lining of the 3rd ventricle and the intermingling of OT dendrites from the right and left side of the ventricle. Since dendritic OT-release stimulates dendritic activity of other OT-neurons, the bilateral proximity of OT-IR dendrites will contribute to the synchronised release of OT by the left and right PVH simultaneously. Abbreviations: *: 3rd Ventricle; f: fornix; AH: anterior hypothalamic nucleus; Pa: paraventricular hypothalamic nucleus (= PVH).




Fig. 3. Figure is reprinted from Ludwig and Pittman [85], with permission from Elsevier. It shows clearly (A) how many OT-IR neurons in the Supraoptic nucleus (SON) extend their dendrites towards the ventral surface of the brain (B). These dendrites (D) are loaded with OT-IR vesicles, sometimes trapped at the moment of dendritic release (small arrows) (C).

release of massive amounts of any peptide along the ventricular wall (either by dendritic release inside the PVH or by axonal release from the varicosities of fibers running in parallel to the ventricular surface [25,26,101–103]) will immediately affect CSF-levels. ‘Short-distance-diffusion’ will occur without delay in the direction of the lower concentrations in the CSF, because there are no intervening barriers. Diffusion will remain ‘CSF-directed’ (‘CSF-petal’) as long as the CSF-levels are the lowest [72]. (The supposed ‘waste-removal’-function of the CSF is fully based on this assumption!) If, however, levels of OT (or any other neuroactive substance) are increased in the CSF, relative to the local extracellular-fluid (ECF)-levels, due to release at some ‘upstream’ location, diffusion will ‘change direction’ and become ‘ECF-directed’ (‘CSF-petal’). (All experiments based on icv-administration are based on this assumption!).

Therefore, increased CSF-levels will sustain and elongate the effects of locally released OT. Because of the free and bidirectional CSF-ECF exchange, all OT-release mechanisms, whether axonal alongside the ventricular walls or via the CSF, are mutually supportive, the main difference being the (short) delay for the CSF-messages.

The exact size and place of the ‘cumulative’ effects’ of CSF-, ECF- and axonally released OT will be dependent on the location and on the local structure of the brain areas involved. The presence of glial cells, astrocyte extensions or fiber layers etc., (all contributing to ‘tortuosity’) may be hindering diffusion in some of the possible directions (leading to ‘anisotropy’) [104–106]. We have been studying tracer injections into the rat brain over the years and we observed frequently that many ‘barriers’ exist that prevent an expected ‘spherical’ diffusion of the injected substances into the surrounding brain tissue (pers.obs, Veening). But if diffusion is hindered in certain directions, it will be enhanced in the other directions, especially towards the barrier-free CSF.

In summary: supra- and sub-ependymal release of OT may contribute to the CSF-OT levels elevated by dendritic release of OT. The many varicose OT-fibers running alongside and terminating in brain areas bordering the ventricular system, suggest a ‘mutual support’-relationship between axonal/terminal released OT and CSF-OT by enhancing and elongating the effects of OT-peaks.

2.5. Synchronisation and cross-talk of OT neurons

Synchronization of OT-neurons seems to be the underlying mechanism for pulsatile release of OT into the blood stream when triggered by suckling [107–109]. However, measurements inside the supraoptic nucleus have shown that synchronisation of dendritic release of OT results in 100 to 1000 times higher levels in the ECF compared to plasma levels [69,110]. In primates OT levels in the CSF may be considerably higher than the peripheral levels and, as mentioned, show an independent rhythm, not observed in plasma levels [40–42]. Circadian CSF rhythms of OT are endogenously generated, leading to five-fold increases in primates [40], synchronized by the daily light–dark cycle [39,40]. These findings suggest that strong positive feedback and synchronizing mechanisms are working during the coordinated dendritic release of OT.

In addition to the remarkable mechanisms involved in the co-activation of OT neurons inside each PVH or supraoptic nucleus, another interesting phenomenon has been observed: cross-talk (‘synchronisation’) between OT-neurons in different brain areas. Most research has been aimed at OT-neurons in the supraoptic nucleus, where it was shown that unilateral activation of specifically OT neurons (and not the vasopressin neurons!) especially in lactating females induce activation in the contralateral supraoptic OT cells [108]. Apparently, all magnocellular OT cells are being synchronized in their activity and OT release, including those in the PVH [111]. Despite the fact that direct reciprocal connections between both supraoptic nuclei have not been observed with the available techniques, there is convincing evidence to exclude diffusion effects [108]. In a careful series of experiments, Moos et al. [109] have shown that ventromedial medullary neurons are involved in the synchronization of OT neurons, including those in the PVH. Interestingly, the ventral medullary brain areas receive messages from the CSF very rapidly because of fast CSF-transport and the ‘vascular pump-mechanisms’ [61–64], and are therefore in an excellent position to provide the appropriate signals for a rapid ‘long-distance-synchronization’. In addition, for the bilateral synchronization of PVH-OT neurons, short axonal collaterals and dendritic extensions across the upper border of the 3rd ventricle [80],

(See Figs. 1, 2) or even volume transmission from one PVH/ventricular wall across the 3rd ventricle to the opposing PVH/ventricular wall (small distance, no intervening barriers) have to be considered alongside the other hypothalamic and extrahypothalamic synchronization sources.

In summary: synchronisation of dendritic OT-release does not only occur between the neurons in a single hypothalamic nucleus, but happens bilaterally for both the PVH and SON, to produce maximal amounts of OT, if necessary.

2.6. OT released into the CSF: are the levels functionally relevant?

Because of the intervening blood-brain-barrier, peripheral OT contributes only a negligible amount (0.002%) to the OT-levels in the CSF [58]. Therefore, OT-levels in the CSF arise from the paraventricular and supraoptic hypothalamic nuclei, directly by dendritic and supraependymal release into the CSF and/or indirectly by terminal release from OT-fibers in other brain areas from where OT may diffuse towards and into the CSF. What do we know about the relative contribution of these sources?

We have to consider that in the magnocellular OT neurons, up to 95% of OT is present in and released from the dendrites [112] which makes this contribution many times more effective in regulating OT-CSF levels than the relative small amounts released by the terminals of the parvocellular neurons projecting to a variety of brain areas. While terminal release, for instance in the central amygdaloid nucleus, may induce a five-fold local increase in extracellular OT [113], it is hard to imagine how such minimal amounts at a considerable distance from the ventricular system could play a role, compared to the 1000-fold increases observed in PVH and SON [83] in the direct vicinity of the ventricular borders. The more so as diffusion over long distances disperses OT in more directions than only towards the CSF. In addition, 'diffusion takes time', which means that the amounts of extracellular OT steadily decrease over time and distance. While the half-life-time for OT in the CSF is about 10 to 20 times longer than in the blood, the ECF-half life-times are probably similar but remain to be determined. Anyway, both time and distance will decrease the amounts of OT reaching the CSF from 'terminal release sites at a distance' to some extent. We propose that the contribution of terminal release of OT elsewhere in the brain (apart from the OT-fibers running along the ventricular system to release their contents along or even purposefully into the CSF) to varying OT-CSF levels will be, at the most, very limited and thereby the idea that varying OT-CSF levels reflect only 'OT-waste-disposal' as utmost improbable.

This raises the question: are the amounts of OT becoming available from the well-coordinated dendritic release in PVH and SON sufficient to explain the varying OT-CSF levels and to induce behavioral/physiological effects? All available evidence suggests: yes, they are! Under resting conditions, basal OT-CSF levels tend to be in the range of 5 to 10 fmol/ml = pg/ml [49,56,114–116], though occasionally higher basal levels have been reported [58,117]. These basal levels tend to increase from about three- up to five-fold higher levels at night in the rat, [39] or at daytime in the human, [41], after osmotic stimulation [116,118], after ejaculation [115] as well as during nursing in guinea pig [49] and sheep [119]. Such increases in OT-levels turned out to be sufficient and have been causally related to effects on nursing [119], learning processes [58] as well as pain-suppression [117]. In the latter publication it was shown that after electrical stimulation of the caudal PVH OT-levels were only increased in the CSF, while after rostral PVH-stimulation increased peripheral OT-levels were induced as well [120], a remarkable case of goal-specific release.

Concerning the amounts of OT necessary to induce such 3 to 5-fold increased OT levels, it was shown by Robinson and Jones [49] that an injection of 100 pg OT into the ventricular system of the guinea pig induced an eightfold increase from about 5 to 40 pg/ml, obviously in the range of the behavioral effects, discussed before. It has been

calculated that the SON contains 3.2 ng OT [112,121]. Adding the magnocellular PVH neurons, the total amount of OT 'bordering' the CSF can be estimated as about 5 or 6 ng OT, of which 95% is available for dendritic release [112]. These amounts are 'huge' compared to the 100 pg necessary for an eightfold increase, which in itself seems to be two times higher than necessary to induce a physiological or behavioral effect. Amounts like 100 pg have, indeed, been shown to become readily available. The concentration of OT in the cisterna magna of the rat increases up to about 300 pg/ml after morphine withdrawal, with the SON as the sole source of this 'subarachnoid OT', because lesioning the PVH was without effect [60,83]. In addition, during acupuncture analgesia in the rat, the SON may release as much as 750 pg/mg protein, completely on its own without support from the PVH, with clearcut analgesic effects [122].

The available evidence suggests that OT neurons in the PVH are perfectly in a situation to work synchronously (by positive feedback as well as by cross-talk between left and right PVH), to release considerable amounts of OT specifically into the rostral 3rd ventricle, without concomitant release in the blood stream, providing potentially an independent message to other brain areas, in addition to the 'parvocellular OT-projections', without peripheral involvement. OT-neurons in the SON may provide a similar message to brain areas at the outer surface of the brain, bordering the subarachnoid space.

In view of these neuroanatomical and functional characteristics of the OT system described above, we prefer to consider the CSF not as a 'drain' by which superfluous amounts of OT are removed, but, following Sowards and Sowards [31], as an actively regulated system carrying variable amounts of OT, as a message to widespread 'downstream' brain areas.

In summary: since systemic OT does not contribute to CSF-OT, all CSF-OT must be derived from PVH and SON by dendritic release, supported by release via supraependymal elements and parvocellular terminals. The amounts of available and experimentally released OT into the CSF seem to be amply sufficient to compose a relevant signal or to 'send a message' to sensitive brain areas.

2.7. Oxytocin receptors and the CSF

Questions about 'sensitive brain areas' focus on the distribution and density of OT receptors, that have been studied by a variety of techniques, in different animals and at different ages [123–131] including the human brain [132,133] and genetically modified lacZ gene- and Venus cDNA-expressing mouse brains [124,131]. In an extensive review, Gimpl and Fahrenholz [134] discussed the OT-receptor system, and they listed about 50 brain areas showing detectable levels of OT-mRNA-expression or the occurrence of OT-binding, in the rat and human brain. A few points, related to the subject of the present review, have to be addressed more specifically.

An OT-message consisting of a single or a few successive peaks in the ventricular OT-levels contains in itself a very limited amount of 'content', hard to reconcile with the many and variable effects of icv-administration of OT, as described in the literature (see below). Necessarily, it has to be conceived that the 'receptive system' (OT-receptors and binding sites) is a variable and plastic part of the communication system. It is hardly surprising, therefore, to see that an extensive set of controlling factors determines the 'reactive state' of the OT-receptive system, with long- and short-term changes occurring both globally and in a site-specific way. It is the variability in the OT-receptive target areas that explains the differentiated behavioral effects of single icv-OT injections [135] (Fig. 4). Recently, it has been shown experimentally that genetic manipulation of the 'receptive system' and viral vector-mediated changes in receptor expression induce impressive long-term behavioral changes [136–140].

As observed by Tribollet et al., [129], the distribution and local densities of OT-receptors are strongly depending on the age of the individual and on existing species differences [129]. Changes in




Fig. 4. Figure is reprinted from: Ostrowski [135], with permission of Elsevier. It shows the distribution of OT-receptor mRNA in the rat brain. The ample occurrence in medial hypothalamic, superficial cortical layers as well as the olfactory regions is immediately clear.

expression of OT-receptors concur with important behavioral changes, while neonatal manipulations induce long-lasting behavioral effects [128,141,142]. Some species-differences are clearly related to social conditions (living monogamously or polygamously) [140,143], but for other differences the functional context is not immediately clear. OT-receptors appear to be under extensive regulatory control, however, exerted by a variety of steroids, circulating in the blood stream. It has been shown that cholesterol [144,145], gonadal steroids [146–149] and corticosteroids [150,151] all play a regulatory role in the density and sensitivity of the OT-receptors. Since steroid levels are constantly changing, due to a large variety of internal as well as external control mechanisms, so will be the case for local OT-receptivity. These steroid effects are not involving the whole population of OT-receptors irrespectively, but seem to be targeting specific brain areas, as shown by the specific variations during the estrous cycle on OT-receptor densities in the ventromedial or the preoptic hypothalamus in the rat [149,152–158]. In this way, a picture of constantly changing patterns of OT-receptivity emerges when several brain areas are considered together. Additional indirect

regulatory effects, like those described for opioids [159,160] may add considerably to the observed variations in OT-receptor densities and distribution. Finally, the ‘common use’ of the AVP1 receptor makes the OT-message partially depending on the state of the AVP-receptive system, adding to local variations in sensitivity for the ‘OT-message’.

Taken together, the ‘OT-receptive system’ impresses as an extremely flexible congregation of brain areas, of which ‘OT-sensitivity’ constantly changes over time, short-term and/or long-term, as well as over location. Apparently, OT-sensitivity should be described as variable patterns. Such a variable patterning of OT-sensitivity over time and location seems to be amply sufficient to understand how a single ‘one-dimensional’ message may have such a variety of effects.

Among the 50 OT-receptive rat brain areas listed by Gimpl and Fahrenholz [134], at least 25 are located at a distance of less than 1 mm from the ventricular walls surrounding the CSF, and when the outer surface of the brain is taken into consideration as well, including the cortical areas, this number increases to about 40, revealing that

80% of the brain areas containing OT-receptors are potentially influenced by volume-transmission via CSF-OT, within a matter of minutes (Fig. 5).

'Mismatch' is a well known phenomenon in the brain, meaning that the distribution of immunoreactive fibers for a given neuroactive brain substance is not congruent with the distribution of its receptors [161,162]. In the case of OT, it is clear that many of the areas containing OT-receptors receive direct OT-projections from mainly the parvocellular part of the PVH. However, some remarkable exceptions seem to be present, suggesting a considerable 'mismatch' between the density of the innervating fibers and the density of OT-binding sites. When comparing the distribution of OT-IR fibers in the rat [65,67] and guinea pig brain [163] with the brain areas showing the highest 'OT-receptivity' [127,129,130,134,135,143,164] several remarkable differences become apparent. Some 'OT-receptive' brain areas hardly receive any OT-innervation, while some other brain areas may contain OT-IR fibers of parvocellular origin, but their close proximity to the ventricular system or subarachnoidal space suggests that interactive neuronal-CSF-OT-effects, as discussed before, most probably occur all the time.

The most interesting brain areas have been indicated in Fig. 5. This figure shows the flow of CSF with the OT 'going with the flow' and 5 areas have been indicated specifically, where functional behavioral effects of OT might be of considerable importance. These brain areas will be discussed in more detail below, in section 4, but comprise medial hypothalamic regions, with important behavioral functions, brainstem areas involved in general and visceral functions, cortical areas and olfactory regions.

In summary: if OT-releasing neurons use the CSF-system for distant messages, the variable, plastic part must be at the receptive side of such a 'messaging-system' and relevant brain areas should be bordering the ventricular and subarachnoid spaces. The great majority of OT-receptive brain areas are found along the ventricular and arachnoid surfaces and some of them do not even contain any OT-IR-fibers ('mismatch'). Density and sensitivity of OT-receptors are extensively controlled by all kinds of steroid hormones and other factors, in a localised and specific way, resulting in a very variable pattern of receptivity of brain areas.

3. Behavioral effects of oxytocin are elicited in the CSF

Functionally, OT and its receptors are involved a wide variety of behaviors like maternal-, sexual-, aggressive-, affiliative-, anxiety-, grooming and feeding behavior, as well as in nociception [134].

The OT-involvement in a particular behavioral system is usually intertwined with other neurotransmitter or endocrine changes, as well as peripheral physiological phenomena. Most of the effects on organs or bodily systems can be understood as adaptive changes, supporting the behavioral effects, mentioned above. Some of the behavioral effects can be induced by (large amounts of) peripheral OT but, generally, peripheral administration induces only peripheral effects, due to the blood-brain-barrier. However, small amounts of icv-OT are remarkably effective to induce behavioral effects, while in some cases, direct neuronal connections of parvocellular OT-neurons seem to be indispensable.

The hypothesis that, besides local release from axons, OT distributed by CSF plays a role in behavior is supported by the many effects found after intracerebroventricular injection of the neuropeptide.

3.1. Social behavior

Social behaviors exist in many forms and have been described as including parental behavior, affiliative behavior, social attachment, sociosexual behavior and monogamy, and all of these share the involvement of OT [140,165–177]. In rodents, sheep and goats, olfactory and other cues, guiding social recognition, are obviously under oxytocinergic control. Several reviews appeared recently and showed the OT-involvement in a series of brain areas, like the olfactory bulb, the medial amygdala, the medial preoptic area, the ventral tegmental area and the nucleus accumbens, [166,177–186]. Repeatedly, the similarities between the effects of disturbed OT systems and clinical syndromes like autism have been pointed out and recently the beneficial clinical effects of OT-treatment on positive social interactions as well as for autism were reported [167,187–191]. Interestingly, underlying the different behavioral reactions to cues to be approached or to be avoided [177], there seems to be a balance




Fig. 5. Figure is based on a midsagittal section of the rat brain, kindly provided by L.W. Swanson. It shows dendritic OT release-sites, from where OT 'goes with the flow'. An 'internal flow' caudalward from the PVH, followed by a rostralward flow along the outside of the brain. The supraoptic nucleus (SON), at the ventral surface of the hypothalamus, mainly contributes to this 'external flow'. The main regions containing OT-receptors and potentially influenced by 'OT-messages' arriving with the CSF, have been indicated and numbered 1 to 5 as follows: 1, 'Behavioral control column' in the hypothalamus; 2 'core' and 'paracore'-brainstem areas; 3, cortical areas; 4, olfactory regions; 5, olfactory sensory neurons. Each of the regions is discussed in the text. OT released in the CSF starts flowing caudally along the dorsal and ventral side of thalamic adhesion, to enter the cerebral aqueduct (AQ) and fourth ventricle; a small part of the CSF may continue flowing caudally through the central canal of the spinal cord, but most of the CSF leaves the ventricular system via the lateral apertures to start flowing through the subarachnoid space, surrounding the brain. This 'external-flow' is indicated here along the dorsal and ventral surface but occurs along all external brain surfaces. The destination of the arachnoid flow is the cribriform plate of the ethmoidal bone, containing the penetrating olfactory fibers, where CSF is released in small lymphatic vessels. The picture suggests that the ventricular bulk-flow of CSF is caudalward and arachnoid flow mainly rostralward. Abbreviations: IVF: interventricular foramen; PVH: paraventricular hypothalamic nucleus; SON: supraoptic hypothalamis nucleus; V3(p,h,m,t,pl): parts of the third ventricle; V4: fourth ventricle; 1–6: circumventricular organs; 5: Pineal organ.

between OT and the pituitary-adrenal axis activity [147,192–196]. In this interaction, OT has stress-reducing effects, which even appear to be transmittable to a conspecific cage mate by olfactory cues [197,198]!

Intranasal administration of a variety of substances has been shown to be effective in inducing physiological, behavioral and clinical effects [199–207]. In humans, intranasal administration of oxytocin causes a substantial increase in trust among humans and decreases anxiety [208–213], thereby greatly increasing the benefits from social interactions. Oxytocin specifically affects an individual's willingness to accept social risks arising through interpersonal interactions [214]. Oxytocin also improves 'mind reading': the ability to infer the mental state of others from physical cues from the eye region [215]. Interestingly, the effects of intranasal administration of OT (as well as many other neuroactive substances) seem to originate not from the 'olfactory system' itself, but from the 'perineuronal spaces' surrounding the olfactory fibers while penetrating the cribriform plate, to reach the olfactory bulb and surrounding liquor space. The identity of and the relationships between these 'perineuronal spaces' and the spaces functioning during the CSF-release into the 'nose-lymphatics', certainly deserve more attention and additional research [199–201,210–212,216,217].

3.2. Sexual behavior

An overwhelming amount of data shows that OT stimulates erection, with both central and peripheral mechanisms involved [218–228]. In the succeeding phases of the copulatory sequence of the male rat, OT significantly reduced mount and intromission latencies only in aging (20-month-old) rats [229], but the reduction in ejaculation latency as well as in the duration of the post-ejaculatory interval were observed also in young vigorous males, most consistently after icv-administration [230,231]. Recently, a case of male anorgasmia was treated successfully by intranasally applied OT [232]. Peripherally, OT stimulates only some specific aspects of the ejaculatory process, like sperm transport [225,233–235]. Generally, peripheral interactions and the integrity of the pelvic nerve seem to be necessary, at least for the effects of systemically administered OT [236].

Interestingly, OT has antidepressant effects in several animal models for depression [237]. Antidepressant drugs, like specific-serotonin-inhibitors (SSRI's) may induce inhibition of masculine behavior [238]. OT may reverse this SSRI-induced inhibition [239,240]. Moreover, oxytocin is released centrally during sexual activity and mating with a receptive female, and this seems to be involved in the reduction of anxiety and the increase in risk-taking behavior in male rats that continues for several hours after copulation [77].

In contrast to the generally stimulating effects of OT on male sexual behavior, inhibitory effects of OT have also been observed, after high doses, especially after icv-administration. These inhibitory effects on penile erection [241] as well as on later phases of masculine behavior [242,243] suggest that larger dosages of OT, administered centrally, may 'create a physiological state, analogous to sexual satiety' [244,245]. Concerning the brain areas involved in these effects on male sexual behavior, the OT-neurons in (and adjoining) the paraventricular hypothalamic nucleus play a crucial role [246], but hippocampal involvement in erection has been described as well [227,228,247].

In addition, OT has been shown to play a facilitatory role in female sexual behavior [230,245,248–253], possibly due to its involvement in the efferent control of the (rodent) clitoris and vagina [254].

Ovarian hormones as well as sexual interaction strongly influence the distribution and intensity of immunoreactivity of OT neurons [255] and OT + Fos-IR neurons have been observed after sexual activity in female rats [256]. Defining the neural substrate for these behavioral effects is complicated, however, although the medial preoptic area and the lateral part of the ventromedial hypothalamic

nucleus seem to be involved [257], precisely the brain areas where OT-receptors are influenced by gonadal hormones [154–158], (see section 2.7)! Part of the OT effects may be indirect by increasing the tolerance of the female rat for tactile stimulation or by actions on affiliative behaviors (see below) associated with sexual behavior [244]. Finally, species differences, even between rodents, may be considerable, because in the prairie vole no facilitatory effects could be demonstrated. Instead, high dosages resulted in immediate and long-lasting inhibition of female sexual behavior, but only after icv and not after systemic administration [244,245].

3.3. Parental behavior

Especially in the peri- and postpartum period, a remarkable set of physiological and behavioral changes is induced, including parturition and lactation [168,258]. To mention just a few of the adaptive changes described: olfactory changes, related to olfactory/social recognition [114,181], changes in brain activation patterns [259,260], morphological and excitatory changes in specific brain areas [92,261] as well as changes in OT release concurrent with receptor expression [262,263]. During parturition, the brain of the newborn is protected by the elevated OT levels [264], while afterwards the anxiolytic effects of OT become more manifest, supporting maternal defence [265]. A recent fMRI study showed the modulating effects of OT on unconditioned fear responses in lactating dams [266]. Apparently, the integrative role of OT around parturition is highly complex and affects several behavioral and physiological levels in order to adjust the physiological mechanisms to the necessary changes in behavior. Many of the changes described have been related to functional changes in the 'OT-receptive' target areas occurring in this time period.

In ewes, intracerebroventricular infusions of oxytocin significantly increased the frequency of some or all of the maternal behaviors scored, while aggressive and negative behaviors significantly decreased in frequency [119]. In vivo microdialysis showed that OT concentrations increased significantly in the region of PVH at birth in multiparous ewes. When OT was retrodialysed bilaterally into the PVH, maternal behavior was induced [267].

3.4. Food intake

In relation to food intake, OT has been shown to affect both peripheral physiological aspects like gastric motility [268,269], gastric acid secretion [270] and insulin secretion [271] as well as central behavioral aspects. OT generally seems to inhibit food intake while OT-knockout mice show enhanced intake of palatable carbohydrate solutions [272]. OT seems to play a special role in linking 'long-term'-satiety signals (leptin) to meal size control. The cells of origin for these inhibitory effects are OT-neurons located in the paraventricular nucleus, and their descending fibers may exert a tonic inhibitory influence via (parts of) the nucleus of the solitary tract. Interestingly, icv-administration of OT induces also strong feeding-related effects, especially via the 4th ventricle, where application of an OT-antagonist was shown to be about 1000-fold more effective than in the 3rd ventricle [273,274]. Taken together, these data are strongly suggesting a conjoined and mutually supporting effort of neuronal and CSF-messaging influencing the brainstem areas that control food intake. Additional effects on more rostral brain areas, like the activation of the ventromedial hypothalamic nucleus [275], previously known as 'satiety center', may further contribute to a subtle differential inhibitory control of food intake.

3.5. Grooming behavior

Grooming behavior can be induced by an icv injection of OT [276–279], even in transgenic 'OT-null'-mice [280]. These effects were

thought to be initiated via the paraventricular hypothalamic nucleus and/or ventral tegmental area [281,282], whereas the nucleus accumbens was associated with inhibition of OT-icv-induced grooming [283].

3.6. Pain

OT has been reported to influence pain control mechanisms [117,122,284–288], modulation of the sympathetic nervous system [289,290], and cerebral hemodynamics [291], suggesting even further going indirect adaptive control functions for OT. Concerning the effects on pain regulating mechanisms, it turned out that only icv-administration of OT was effective in elevating the pain threshold, which could be blocked by an OT-antagonist, while peripheral administration was completely ineffective [122]. Increased OT-levels in the CSF, induced by central release of OT after PVH-stimulation, were directly involved in and causally related to the pain suppression [117].

In summary, icv-injections of OT induce many different behavioral as well as pain-reducing effects. Apparently, the state of the 'OT-receptive-system' determines the effect of an experimental 'CSF-OT-peak'. Systemic administration of OT frequently induced no behavioral effects at all and was completely ineffective in pain suppression.

4. CSF-oxytocin may affect various parts of the CNS

The multiple and variable behavioral effects of icv-injections of OT suggest a specific role for the CSF in distributing OT-messages to distant destinations. Fig. 5 summarizes how OT, released from the PVH into the ventricular system and from the SON into the subarachnoidal space, reaches many OT-receptive parts of the brain by 'going with the flow'. Some of these brain areas deserve special attention, because of their strong involvement in a variety of physiological, behavioral and cognitive functions. These areas have been numbered 1 to 5 in Fig. 5, and will be discussed successively.

4.1. OT- effects on hypothalamic 'Behavioral Control Column' (Nr 1 in Fig. 5, Fig. 6)

Over the last years, Larry Swanson has developed the concept of a 'behavioral control column' [1,292]. This column consists of a longitudinal series of nuclei extending from the most rostral (preoptic) to the most caudal (pre-mammillary and mammillary) parts of the medial hypothalamus, including the anterior and ventromedial nuclei. Nuclei of this hypothalamic 'behavioral control column' turn out to play a keyrole in the organisation of behavior. (Fig. 6). Many neurons of this 'column' are steroid sensitive and Pfaff and co-workers have shown over the years that circulating gonadal hormones directly affect OT-receptivity in the lateral part of the ventromedial hypothalamic nucleus, involved in the lordosis-circuitry of the female rat [154–156,158,256,257,293–300]. This impressive line of research shows very clearly how steroid hormones influence OT-receptivity, allowing the gonadal endocrine systems regulatory control over a range of other behaviors. As an example: female rats readily switch their response to a male from a soliciting proceptive attitude when in estrous to an aggressive response when in non-estrous, showing how different hormonal states induce radically different reactions towards the same approaching stimulus. In this kind of behavioral reactions, the 'behavioral control column' plays a keyrole together with some closely (and mostly reciprocally) connected parts of the bed nucleus of the stria terminalis, the amygdaloid complex and maybe other medial and cortical parts of the 'limbic system' [30].

As mentioned before, OT and its receptors induce a complex set of behavioral changes, especially peri- and postpartum, involving effects on maternal, sexual and affiliative behavior, female aggression,

anxiety, nociception and social memory [134]. One or more of the hypothalamic nuclei of the 'behavioral control column' are always involved in these behavioral changes, as supported by numerous Fos- and other studies from our group [301–306] as well as from many others. Each of the nuclei of the 'behavioral control column' is located closely to the ventricular walls (in rats (much) less than a mm) and contains neurons provided (sometimes densely) with OT-receptors [130].

From these facts we propose that some major target areas of CSF-OT are located in the medial-hypothalamic 'behavioral control column' as defined by Swanson. The strong effect of icv-OT on ventromedial hypothalamic neuronal activity [275] is a nice demonstration of such hypothalamic targets. Suppression of aggressive reactions, normally obtained from the anterior hypothalamic nucleus, or the adjoining 'hypothalamic attack areas' [307–311] could obviously be another OT-effect realised in the 'behavioral control column'. In addition to possible CSF-effects, some rostral (preoptic) and some caudal (pre)mamillary nuclei may receive a direct OT-innervation as well, supporting the idea of mutually supporting messages.

N.B. Our conclusion regarding the importance of these paracrine OT-effects are neither meant to indicate that these paracrine effects are the only ones influencing these target areas, nor that the target areas are restricted to this behavioral control column!

4.2. OT- effects on brainstem ('core'- and 'paracore'-) areas (Fig. 5, nr 2)

Since there is no reason to assume that the caudal flow of CSF-OT becomes ineffective at the entrance of the cerebral aqueduct, the receptive parts of the periaqueductal gray (PAG), raphe nuclei, locus coeruleus and the solitary tract nucleus [30,31,134,312,313] may all add their contribution to the behavioral changes observed. Especially the PAG seems to be easily accessible for all kinds of substances passing the cerebral aqueduct, as apparent from all studies concerning the penetration depth of substances added to the CSF [161,314–316] as well as those concerning GnRH effects on the PAG [317,318].

However, brain areas more remote from the ventricular or subarachnoidal space may also have easy access to (the contents of) the CSF as well, and tanycytes seem to play a crucial role here. Tanycytes are specialised bipolar ependymal cells contacting the CSF [319–324], especially in the mediobasal hypothalamus. However, brainstem areas like locus coeruleus, raphe nuclei and the paraventricular nucleus in the ventral brainstem are also provided by tanycytes [325–329]. Very recently it was shown that the raphe nuclei may be reached by CSF-influences via such specialised tanycytes [330] and that the locus coeruleus accumulates nerve growth factor, about 50% of it not by neuronal transport mechanisms but obtained from the CSF [331]. Assuming that these findings form part of an increasing series of similar findings, it is interesting to recognize that the raphe nuclei and the locus coeruleus form part of the 'medial and lateral paracore areas' respectively, as defined by Nieuwenhuys [7,20,22]. These 'paracore areas' extending through the brainstem are characterized by their content of numerous unmyelinated peptidergic fibers, apparently involved in non-synaptic communication and subserving the physiological adaptations necessary for appropriate emotional and behavioral reactions [7]. We hypothesize that CSF-messages become effective in these brainstem areas in cooperation with the many unmyelinated fibers, releasing neuropeptides in many brainstem areas, frequently running close and parallel to the ventricular system, especially the cerebral aqueduct.

4.3. OT- effects on cortical areas (Fig. 5, nr 3)

Another specific and important part of the brain influenced by the OT-CSF-flow, is the cortical mantle, covering most of the telencephalon. Interestingly, a direct cortical innervation by OT-fibers seems to




Fig. 6. Figure is a horizontal reconstruction of the rat brain, kindly provided by L.W. Swanson. The concept of a 'Behavioral Control Column' [1,292,367] is represented by the series of hypothalamic nuclei, extending from the rostral preoptic part to the caudal premammillary region. While the number of OT-IR fibers traversing the medial hypothalamus is very limited, the presence of OT-receptors is generally abundant, as indicated by the orange arrows in the lower part of the figure, that are based on Vaccari [130]. As discussed in the text, this situation suggests that the 'Behavioral Control Column' can be considered as one of the main target regions for OT-messages flowing with the CSF. Abbreviations: MPN: medial preoptic nucleus; AHN: anterior hypothalamic nucleus; PVHd: paraventricular hypothalamic nucleus; VMH: ventromedial hypothalamic nucleus; PMv & PMd: ventral and dorsal premammillary nuclei; VTA: ventral tegmental area.

be virtually or totally lacking in many cortical areas. Nevertheless, quite a few cortical areas are OT-receptive, with the highest densities of OT-receptors located in cortical layers 2 and 3 [127,130,134,135]. Since cortical neurons located in layers 2 and 3 are mostly involved in the intracortical connectivity, the following picture emerges: OT arriving via the CSF, has easy and rapid access to cortical and deeper brain areas via the perivascular spaces. OT may spread over the whole extension of the cortical mantle, to influence OT-receptive neurons in layers 2 and 3, not only to influence cortical development [332] but also to induce changes in local or overall cortical activity or functional 'connectivity'. Experimental data, confirming that OT affects cortical activity, including attenuation of memory and learning [333] are not difficult to find, despite the fact that these cortical effects of OT received considerably less attention, so far, than the subcortical behavioral effects [334–337].

Recent findings show the key role of OT in phenomena like 'affective evaluation' and 'trust' [338–340]. In the human brain, 'Von Economo neurons' play an important role in autism, intuition and social behavior [341–343]. The possibility of effective treatment regimens by intranasal OT-application offers exciting new opportunities [215,338,344–346] and it will certainly be possible to define the role of cortical OT much more specifically, in the near future.

4.4. OT- effects on olfactory areas (Fig. 5, nr 4)

If most of the CSF and OT is drained into the olfactory lymphatics, it means that the olfactory bulbs as well as all adjoining superficial olfactory areas are fully exposed to the surrounding CSF-OT, which makes it less surprising that electrical stimulation of the paraventricular hypothalamic nucleus (PVH) induces the same inhibitory effect on the mitral cells in the olfactory bulb as microinjection of oxytocin in the lateral ventricle (icv) [347]. While there are some OT-fibers directly innervating the bulbs and other olfactory areas,

unilateral transections between PVH and olfactory bulb failed to block the stimulation effects and the authors suggested explicitly that the CSF was the way of communication [347]. Anyway, the density of OT-receptors in the olfactory areas is generally high [130,134,135,348] and easily accessible for 'CSF-OT'.

Taking into account the location of the easily accessible olfactory bulbs, completely surrounded by CSF, it is no longer surprising that OT is very effective in influencing in 'olfaction-related-behaviors' and or can be used as a target for intranasal-OT sprays [344,346].

4.5. OT- effects on olfactory sensory neurons (OSNs) (Fig. 5, nr 5)

Drainage of CSF via the numerous small lymphatic vessels of the nose, opens the possibility that eventually even the olfactory sensory neurons (OSNs) themselves could be affected by the temporarily elevated OT-levels, while at the same time linking the brain to the lymphatic system [349]. Such a hypothetical effect of OT on the OSNs would serve the function of adapting olfactory sensitivity to the hormonal and behavioral state of the animal. This idea seems attractive for a number of reasons: 1) changes in the motivational state are known to induce temporary adaptive changes in the somatosensory, gustatory, visual and auditory sensations, and there is no reason to assume that the olfactory sense would be an exception to what seems to be a general rule: motivational changes induce sensory adaptations. 2) gonadal hormones have been shown to influence olfactory sensations extensively (factors like estrous vs anestrous, or effects of castration [350–352]). 3) direct effects of neuropeptides on olfaction are harder to find, but in 'behavioral states' like hunger vs satiety, two neuropeptides play a prominent role: orexin (of hypothalamic origin and stimulating food intake) and leptin (of peripheral origin and inhibiting food intake) [353–357]. Note that these peptides not only affect hypothalamic/limbic mechanisms regulating food-intake but also show direct effects on

the olfactory sensations themselves, just like the motivational states hunger and satiety have been shown to induce changes in olfactory sensitivity [358–360]. 4) for OT to exert direct effects on the OSNs, the latter necessarily would need to be in the possession of the appropriate receptors. This information is hard to find in the literature, which reflects a general lack of knowledge about possible modulating effects of neuropeptides on OSNs. However, Levasseur et al. [361] have studied cultured OSNs and they showed not only the presence of several types of receptors on OSNs, but also direct effects of vasopressin and OT on OSNs. Especially the effect of OT, at this cellular level, turned out to be strong!

Several other neuropeptides have been shown to affect OSN sensitivity: in addition to orexin and leptin, mentioned above, substance P [362], natriuretic peptides [363] and GnRH [364]. These findings strongly suggest that it is probably the rule, and not an exception, that neuropeptides involved in motivational changes, concurrently modulate olfactory sensitivity into a state optimally adapted to the current behavioral state.

Finally we would like to ask attention again for the puzzling findings of Agren et al., who showed that the stress-reducing effects of OT are apparently transmitted to a conspecific cage mate by olfactory cues [197,198]. Could it be that the OT in the nose-lymphatics and nasal epithelium is playing a role here?

There is much experimental work to be done in this field of the functional relations between olfactory bulbs, the surrounding CSF, the olfactory fibers and adjoining lymph vessels and the possible direct modulating effects of neuropeptides as well as other substances on OSNs. Since each of these structures may also be involved in the effects of 'intranasal administration' of drugs and neuropeptides, which treatment is growing more and more important as an alternative route for drug administration, to circumvent the blood-brain-barrier, it is to be expected that considerably more experimental data and more conclusive evidence will come available over the next few years.

In summary: The most interesting brain areas, potentially targeted by OT-messages via the CSF, comprise the medial hypothalamus and several brainstem, cortical and olfactory regions. For each of these regions, the available evidence that OT might be involved in functional and/or behavioral changes is presented. Apparently, there remains a lot of work to be done.

5. Conclusions and future directions

In the previous sections we have stated that the CSF is used for communication in the central nervous system, that oxytocin is released into the CSF, that oxytocin in the CSF contains a meaningful message and reaches many brain areas that contain functional oxytocin receptors and that oxytocin in the CSF has multiple marked behavioral effects.

The question that remains is: what is the functional significance and the specific contribution of this relatively uncontrolled and broad-spectrum oxytocin release system, in addition to the targeted axonal release from parvocellular OT-neurons and the peripheral release of the magnocellular OT-neurons? We support and propose the view that broadcasting messages via the CSF is an appropriate way to put several parts of the CNS into a 'behavioral state', subserving the performance of specific behaviors or groups of behavioral elements sharing a common motivational system including the temporary suppression of specific behavioral or physiological reactions. These 'CSF-messages' have to be considered as a partially independent signal, while at the same time constantly supporting, sustaining and in mutual exchange with the ECF-contents released by axon terminals all over the brain, but mostly in brain areas adjoining the CSF. This signaling pathway has been described recently as an 'undervaluated pathway of neuroendocrine signaling into the brain' [365]. The

presence of nanostructures, from various brain sources, appears to support such signaling functions in the human CSF [366].

Sewards and Swards [31] proposed that the presence of elevated CSF- levels of corticotropin releasing factor (CRF) and vasopressin generate fear and 'power-dominance' drive motivation, respectively. They hypothesized that the elevated neuropeptide levels in the CSF are detected and transduced into neuronal activities by, predominantly, hypothalamic neurons in the vicinity of the third ventricle. The purpose of this system was proposed to be maintaining a state of fear or anger and consequent vigilant or aggressive behavior after the initial fear- or anger-inducing stimulus itself is no longer perceptible [31]. In other words, CSF-messages prolong the duration of a motivated state. Interestingly, important destination areas for this transmission (the medial hypothalamus, periaqueductal gray, midline thalamus and medial) prefrontal cortex are all bordering the CSF-liquor system, at the inner- or outer surface of the brain. In addition, each of these destination areas is extensively connected to other parts of the limbic system and functionally involved in a range of differently motivated behaviors [31].

We propose that a similar (but compared to CRF opposite) role is being played by a CSF-OT-message-system. OT-release into the CSF upon stimuli like parturition, suckling or hand-milking, vagino-cervical stimulation or mating, can lead to a number of behavioral effects: facilitation of sexual behavior, inducing partner preference or pair bonds, facilitation of maternal behavior, decrease of aggressive behavior, increase of physical contact, increase of proximity and coherence in a group of huddling pups, increase in trust, improvement of social 'mind-reading' and elevation of the pain threshold, amongst others.

Although the complete scope of behaviors related to CSF-oxytocin release may not yet be complete, the general pattern that emerges is one of an increase in social behavior, contacts and 'trust', combined with a decrease in stress and anxiety'.

In summary: We propose that OT-messages are broadcasted via the CSF in order to maintain for a prolonged period of time a 'pro-social' behavioral state, allowing and increasing bodily contacts, while at the same time modulating a variety of other behaviors, depending on the sensitivity (or 'OT-receptive state') of numerous brain areas involved.

Considering all available evidence we propose that:

- OT levels in the CSF are being regulated actively;
- OT levels in the CSF contain a meaningful message for brain areas in the appropriate receptive state;
- OT-messages via the CSF are effective at an appropriate timescale to induce behavioral changes as well as to support and sustain the effects of OT released by axonal terminals in many 'behavior-relevant' brain areas bordering the ventricular and subarachnoid spaces;
- the 'behavioral control column' in the hypothalamus, brainstem areas ('core' and 'paracore' as well as ventral superficial) and the cortical and olfactory areas bordering the subarachnoid space and most probably including the olfactory sensory neurons, can be considered as the main target areas for the CSF-OT messaging system.

References

- [1] Swanson LW. Brain architecture, understanding the basic plan. New York: Oxford University Press; 2003. p. 263.
- [2] Hinde RA. Animal behaviour; a synthesis of ethology and comparative psychology. second ed. New York: McGRAW-Hill Book Cy; 1970.
- [3] Mogenson GJ. The neurobiology of behavior: an introduction. Hillsdale: Lawrence Erlbaum Associates; 1977. p. 334.
- [4] Barendregt H. Buddhist Phenomenology (Part 1). Conference on Topics and Perspectives of Contemporary Logic and Philosophy of Science. Cesena, Italy: Clueb, Bologna; 1988.
- [5] Frijda N. The laws of emotions. London: Psychology Press; 2007.

- [6] Panksepp J. Affective Neuroscience. The foundations of human and animal emotions. New York: Oxford University Press; 1998. p. 466.
- [7] Nieuwenhuys R, Veening JG, van Domburg P. Core and paracores; some new chemoarchitectural entities in the mammalian neuraxis. *Acta Morphol Neerl Scand* 1988;26:131–63.
- [8] Veening JG. *Eur J Morphol* 1992;30:53–66.
- [9] Agnati LF, Fuxe K. Volume transmission as a key feature of information handling in the central nervous system possible new interpretative value of the Turing's B-type machine. *Prog Brain Res* 2000;125:3–19.
- [10] Agnati LF, Fuxe K, Zoli M, Ozini I, Toffano G, Ferraguti F. A correlation analysis of the regional distribution of central enkephalin and beta-endorphin immunoreactive terminals and of opiate receptors in adult and old male rats. Evidence for the existence of two main types of communication in the central nervous system: the volume transmission and the wiring transmission. *Acta Physiol Scand* 1986;128:201–7.
- [11] Agnati LF, Guidolin D, Guescini M, Genedani S, Fuxe K. Understanding wiring and volume transmission. *Brain Res Rev* 2010;64:137–59.
- [12] Fuxe K, Dahlstrom AB, Jonsson G, Marcellino D, Guescini M, Dam M, Manger P, Agnati L. The discovery of central monoamine neurons gave volume transmission to the wired brain. *Prog Neurobiol* 2010;90:82–100.
- [13] Zoli M, Jansson A, Sykova E, Agnati LF, Fuxe K. Volume transmission in the CNS and its relevance for neuropsychopharmacology. *Trends Pharmacol Sci* 1999;20:142–50.
- [14] Agnati LF, Cortelli P, Biagini G, Bjelke B, Fuxe K. Different classes of volume transmission signals exist in the central nervous system and are affected by metabolic signals, temperature gradients and pressure waves. *NeuroReport* 1994;6:9–12.
- [15] Bjelke B, England R, Nicholson C, Rice ME, Lindberg J, Zoli M, Agnati LF, Fuxe K. Long distance pathways of diffusion for dextran along fibre bundles in brain. Relevance for volume transmission. *NeuroReport* 1995;6:1005–9.
- [16] Bjelke B, Fuxe K. Intraventricular beta-endorphin accumulates in DARPP-32 immunoreactive tanycytes. *NeuroReport* 1993;5:265–8.
- [17] Bjelke B, Fuxe K, Agnati LF. Increased diffusion of prolactin-like material into the brain neuropil from homologous adenohipophyseal transplants in the rat neostriatum after a 6-OH-dopamine induced degeneration of the mesostriatal dopamine neurons. *Neurosci Lett* 1989;107:33–8.
- [18] Nicholson C. Signals that go with the flow. *Trends Neurosci* 1999;22:143–5.
- [19] Nicholson C, Chen KC, Hrabetova S, Tao L. Diffusion of molecules in brain extracellular space: theory and experiment. *Prog Brain Res* 2000;125:129–54.
- [20] Nieuwenhuys R. Chemoarchitecture of the Brain. Berlin: Springer-Verlag; 1985. p. 1–246.
- [21] Nieuwenhuys R. Comparative aspects of volume transmission, with sidelight on other forms of intercellular communication. *Prog Brain Res* 2000;125:49–126.
- [22] Nieuwenhuys R, Voogd, van Huijzen. *The Human Central Nervous System*. 4th ed. Berlin: Springer; 2008. p. 970.
- [23] Veening JG, Barendregt HP. The regulation of brain states by neuroactive substances distributed via the cerebrospinal fluid; a review. *Cerebrospinal Fluid Res* 2010;7:1.
- [24] Buma P. Characterization of luteinizing hormone-releasing hormone fibres in the mesencephalic central grey substance of the rat. *Neuroendocrinology* 1989;49:623–30.
- [25] Buma P, Veening J, Hafmans T, Joosten H, Nieuwenhuys R. Ultrastructure of the periaqueductal grey matter of the rat: an electron microscopical and horseradish peroxidase study. *J Comp Neurol* 1992;319:519–35.
- [26] Veening JG, Buma P, ter Horst GJ, Roeling TAP, Luiten PGM, Nieuwenhuys R. Hypothalamic projections to the PAG in the rat: topographical, immunoelectromicroscopical and functional aspects. In: Depaulis A, Bandler R, editors. *The Midbrain Periaqueductal Gray Matter*. New York: Plenum Press; 1991 pp. 387–415.
- [27] Mouton LJ, Kerstens L, Van der Want J, Holstege G. Dorsal border periaqueductal gray neurons project to the area directly adjacent to the central canal ependyma of the C4-T8 spinal cord in the cat. *Exp Brain Res* 1996;112:11–23.
- [28] Cushing H, Goetsch E. Concerning the secretion of the infundibular lobe of the pituitary body and its presence in the cerebrospinal fluid. *Am J Physiol* 1910;27:60–86.
- [29] Friedman GS, Friedman MH. An examination of cerebrospinal fluid for oxytocinergic activity as tested by the rabbit uterine fistula. *Am J Physiol* 1932;103:244–54.
- [30] Swards TV, Swards MA. Representations of motivational drives in mesial cortex, medial thalamus, hypothalamus and midbrain. *Brain Res Bull* 2003;61:25–49.
- [31] Swards TV, Swards MA. Fear and power-dominance motivation: proposed contributions of peptide hormones present in cerebrospinal fluid and plasma. *Neurosci Biobehav Rev* 2003;27:247–67.
- [32] Malpoux B, Daveau A, Maurice-Mandon F, Duarte G, Chemineau P. Evidence that melatonin acts in the premammillary hypothalamic area to control reproduction in the ewe: presence of binding sites and stimulation of luteinizing hormone secretion by in situ microimplant delivery. *Endocrinology* 1998;139:1508–16.
- [33] Skinner DC, Caraty A, Evans NP. Does gonadotropin-releasing hormone in the cerebrospinal fluid modulate luteinizing hormone release? *Neuroendocrinology* 1998;67:37–44.
- [34] Skinner DC, Caraty A, Malpoux B, Evans NP. Simultaneous measurement of gonadotropin-releasing hormone in the third ventricular cerebrospinal fluid and hypophyseal portal blood of the ewe. *Endocrinology* 1997;138:4699–704.
- [35] Skinner DC, Malpoux B. High melatonin concentrations in third ventricular cerebrospinal fluid are not due to Galen vein blood recirculating through the choroid plexus. *Endocrinology* 1999;140:4399–405.
- [36] Skinner DC, Malpoux B, Delaleu B, Caraty A. Luteinizing hormone (LH)-releasing hormone in third ventricular cerebrospinal fluid of the ewe: correlation with LH pulses and the LH surge. *Endocrinology* 1995;136:3230–7.
- [37] Tricoire H, Moller M, Chemineau P, Malpoux B. Origin of cerebrospinal fluid melatonin and possible function in the integration of photoperiod. *Reprod Suppl* 2003;61:311–21.
- [38] Du Vigneaud V, Ressler C, Trippett S. The sequence of amino acids in oxytocin, with a proposal for the structure of oxytocin. *J Biol Chem* 1953;205:949–57.
- [39] Devarajan K, Rusak B. Oxytocin levels in the plasma and cerebrospinal fluid of male rats: effects of circadian phase, light and stress. *Neurosci Lett* 2004;367:144–7.
- [40] Artman HG, Reppert SM, Perlow MJ, Swaminathan S, Oddie TH, Fisher DA. Characterization of the daily oxytocin rhythm in primate cerebrospinal fluid. *J Neurosci* 1982;2:598–603.
- [41] Amico JA, Tenicela R, Johnston J, Robinson AG. A time-dependent peak of oxytocin exists in cerebrospinal fluid but not in plasma of humans. *J Clin Endocrinol Metab* 1983;57:947–51.
- [42] Amico JA, Challinor SM, Cameron JL. Pattern of oxytocin concentrations in the plasma and cerebrospinal fluid of lactating rhesus monkeys (*Macaca mulatta*): evidence for functionally independent oxytocinergic pathways in primates. *J Clin Endocrinol Metab* 1990;71:1531–5.
- [43] Seckl JR, Lightman SL. Diurnal rhythm of vasopressin but not of oxytocin in the cerebrospinal fluid of the goat: lack of association with plasma cortisol rhythm. *J Endocrinol* 1987;114:477–82.
- [44] Devarajan K, Marchant EG, Rusak B. Circadian and light regulation of oxytocin and parvalbumin protein levels in the ciliated ependymal layer of the third ventricle in the C57 mouse. *Neuroscience* 2005;134:539–47.
- [45] Takeda S, Kuwabara Y, Mizuno M. Effects of pregnancy and labor on oxytocin levels in human plasma and cerebrospinal fluid. *Endocrinol Jpn* 1985;32:875–80.
- [46] Kendrick KM, Keverne EB, Hinton MR, Goode JA. Cerebrospinal fluid and plasma concentrations of oxytocin and vasopressin during parturition and vaginocervical stimulation in the sheep. *Brain Res Bull* 1991;26:803–7.
- [47] Neumann I, Russell JA, Wolff B, Landgraf R. Naloxone increases the release of oxytocin, but not vasopressin, within limbic brain areas of conscious parturient rats: a push-pull perfusion study. *Neuroendocrinology* 1991;54:545–51.
- [48] Seckl JR, Lightman SL. Potentiation of lactation-induced oxytocin secretion by intracerebroventricular oxytocin in the conscious goat. *J Endocrinol* 1988;116:273–7.
- [49] Robinson IC, Jones PM. Oxytocin and neurophysin in plasma and CSF during suckling in the guinea-pig. *Neuroendocrinology* 1982;34:59–63.
- [50] van Wimersma Greidanus TB, Janssen S, Frankhuijzen-Sierevogel JC, Maigret C, van de Heijning BJ. Effect of central administration of the kappa-opiate receptor agonist U 69,593 on neurohypophysial hormone levels in blood and cerebrospinal fluid. *Neuropeptides* 1996;30:452–5.
- [51] Kendrick KM, Keverne EB. Effects of intracerebroventricular infusions of naltrexone and phentolamine on central and peripheral oxytocin release and on maternal behaviour induced by vaginocervical stimulation in the ewe. *Brain Res* 1989;505:329–32.
- [52] Williams JR, Insel TR, Harbaugh CR, Carter CS. Oxytocin administered centrally facilitates formation of a partner preference in female prairie voles (*Microtus ochrogaster*). *J Neuroendocrinol* 1994;6:247–50.
- [53] Ivanyi T, Wiegant VM, de Wied D. Differential effects of emotional and physical stress on the central and peripheral secretion of neurohypophysial hormones in male rats. *Life Sci* 1991;48:1309–16.
- [54] Seckl JR, Lightman SL. Effect of naloxone on oxytocin and vasopressin release during vaginocervical stimulation in the goat. *J Endocrinol* 1987;115:317–22.
- [55] Jones PM, Robinson IC, Harris MC. Release of oxytocin into blood and cerebrospinal fluid by electrical stimulation of the hypothalamus or neural lobe in the rat. *Neuroendocrinology* 1983;37:454–8.
- [56] Hattori T, Sundberg DK, Morris M. Central and systemic oxytocin release: a study of the paraventricular nucleus by in vivo microdialysis. *Brain Res Bull* 1992;28:257–63.
- [57] Dogterom J, Van Wimersma Greidanus TB, Swabb DF. Evidence for the release of vasopressin and oxytocin into cerebrospinal fluid: measurements in plasma and CSF of intact and hypophysectomized rats. *Neuroendocrinology* 1977;24:108–18.
- [58] Mens WB, Witter A, van Wimersma Greidanus TB. Penetration of neurohypophysial hormones from plasma into cerebrospinal fluid (CSF): half-times of disappearance of these neuropeptides from CSF. *Brain Res* 1983;262:143–9.
- [59] Jones PM, Robinson IC. Differential clearance of neurophysin and neurohypophysial peptides from the cerebrospinal fluid in conscious guinea pigs. *Neuroendocrinology* 1982;34:297–302.
- [60] Robinson IC, Coombes JE. Neurohypophysial peptides in cerebrospinal fluid: an update. *Ann N Y Acad Sci* 1993;689:269–84.
- [61] Hadaczek P, Yamashita Y, Mirek H, Tamas L, Bohn MC, Noble C, Park JW, Bankiewicz K. The "perivascular pump" driven by arterial pulsation is a powerful mechanism for the distribution of therapeutic molecules within the brain. *Mol Ther* 2006;14:69–78.
- [62] Rennels ML, Blaumanis OR, Grady PA. Rapid solute transport throughout the brain via paravascular fluid pathways. *Adv Neurol* 1990;52:431–9.
- [63] Rennels ML, Gregory TF, Blaumanis OR, Fujimoto K, Grady PA. Evidence for a 'paravascular' fluid circulation in the mammalian central nervous system, provided by the rapid distribution of tracer protein throughout the brain from the subarachnoid space. *Brain Res* 1985;326:47–63.
- [64] Stoodley MA, Brown SA, Brown CJ, Jones NR. Arterial pulsation-dependent perivascular cerebrospinal fluid flow into the central canal in the sheep spinal cord. *J Neurosurg* 1997;86:686–93.

- [65] Buijs RM. Immunocytochemical demonstration of vasopressin and oxytocin in the rat brain by light and electron microscopy. *J Histochem Cytochem* 1980;28:357–60.
- [66] Swanson LW, Sawchenko PE. Hypothalamic integration: organization of the paraventricular and supraoptic nuclei. *Annu Rev Neurosci* 1983;6:269–324.
- [67] Buijs RM. Intra- and extrahypothalamic vasopressin and oxytocin pathways in the rat. Pathways to the limbic system, medulla oblongata and spinal cord. *Cell Tissue Res* 1978;192:423–35.
- [68] Rousselot P, Papadopoulos G, Merighi A, Poulain DA, Theodosis DT. Oxytocinergic innervation of the rat spinal cord. An electron microscopic study. *Brain Res* 1990;529:178–84.
- [69] Landgraf R, Neumann ID. Vasopressin and oxytocin release within the brain: a dynamic concept of multiple and variable modes of neuropeptide communication. *Front Neuroendocrinol* 2004;25:150–76.
- [70] Krisch, B., ed. Ultrastructure of regulatory neuroendocrine neurons and functionally related structures. Morphology of Hypothalamus and its connections., ed. Ganten, D., Pfaff, D. Vol. 7. 1986, Springer-Verlag: Berlin. 251–290
- [71] Vigh B, Manzano e Silva MJ, Frank CL, David C, Czirok SJ, Vincze C, Racz G, Lukats A, Szel A. The circumventricular organs of the brain: do they represent a cerebrospinal fluid-dependent regulatory system? *Med Hypotheses Res* 2004;1:77–100.
- [72] Hoistad M, Samskog J, Jacobsen KX, Olsson A, Hansson HA, Brodin E, Fuxe K. Detection of beta-endorphin in the cerebrospinal fluid after intrastriatal microinjection into the rat brain. *Brain Res* 2005;1041:167–80.
- [73] Abbott NJ. Evidence for bulk flow of brain interstitial fluid: significance for physiology and pathology. *Neurochem Int* 2004;45:545–52.
- [74] Weller RO, Djuanda E, Yow HY, Carare RO. Lymphatic drainage of the brain and the pathophysiology of neurological disease. *Acta Neuropathol* 2009;117:1–14.
- [75] Brunton PJ, Sabatier N, Leng G, Russell JA. Suppressed oxytocin neuron responses to immune challenge in late pregnant rats: a role for endogenous opioids. *Eur J Neurosci* 2006;23:1241–7.
- [76] Leng G, Ludwig M. Jacques Benoit Lecture. Information processing in the hypothalamus: peptides and analogue computation. *J Neuroendocrinol* 2006;18:379–92.
- [77] Waldherr M, Neumann ID. Centrally released oxytocin mediates mating-induced anxiolysis in male rats. *Proc Natl Acad Sci U S A* 2007;104:16681–4.
- [78] Morris JF, Pow DV. New anatomical insights into the inputs and outputs from hypothalamic magnocellular neurons. *Ann N Y Acad Sci* 1993;689:16–33.
- [79] Rhodes CH, Morrell JI, Pfaff DW. Immunohistochemical analysis of magnocellular elements in rat hypothalamus: distribution and numbers of cells containing neurophysin, oxytocin, and vasopressin. *J Comp Neurol* 1981;198:45–64.
- [80] Armstrong WE, Warach S, Hatton GI, McNeill TH. Subnuclei in the rat hypothalamic paraventricular nucleus: a cytoarchitectural, horseradish peroxidase and immunocytochemical analysis. *Neuroscience* 1980;5:1931–58.
- [81] Rho JH, Swanson LW. A morphometric analysis of functionally defined subpopulations of neurons in the paraventricular nucleus of the rat with observations on the effects of colchicine. *J Neurosci* 1989;9:1375–88.
- [82] Pow DV, Morris JF. Dendrites of hypothalamic magnocellular neurons release neurohypophysial peptides by exocytosis. *Neuroscience* 1989;32:435–9.
- [83] Ludwig M, Leng G. Dendritic peptide release and peptide-dependent behaviours. *Nat Rev Neurosci* 2006;7:126–36.
- [84] Morris JF, Ludwig M. Magnocellular dendrites: prototypic receiver/transmitters. *J Neuroendocrinol* 2004;16:403–8.
- [85] Ludwig M, Pittman QJ. Talking back: dendritic neurotransmitter release. *Trends Neurosci* 2003;26:255–61.
- [86] Yamashita H, Okuya S, Inenaga K, Kasai M, Uesugi S, Kannan H, Kaneko T. Oxytocin predominantly excites putative oxytocin neurons in the rat supraoptic nucleus in vitro. *Brain Res* 1987;416:364–8.
- [87] Moos F, Richard P. Paraventricular and supraoptic bursting oxytocin cells in rat are locally regulated by oxytocin and functionally related. *J Physiol* 1989;408:1–18.
- [88] de Kock CP, Wierda KD, Bosman LW, Min R, Kokksma JJ, Mansvelter HD, Verhage M, Brussaard AB. Somatodendritic secretion in oxytocin neurons is upregulated during the female reproductive cycle. *J Neurosci* 2003;23:2726–34.
- [89] Wang H, Ward AR, Morris JF. Oestradiol acutely stimulates exocytosis of oxytocin and vasopressin from dendrites and somata of hypothalamic magnocellular neurons. *Neuroscience* 1995;68:1179–88.
- [90] Warembourg M, Poulain P. Presence of estrogen receptor immunoreactivity in the oxytocin-containing magnocellular neurons projecting to the neurohypophysis in the guinea-pig. *Neuroscience* 1991;40:41–53.
- [91] Theodosis DT, Poulain DA. Neuronal-glia and synaptic plasticity in the adult rat paraventricular nucleus. *Brain Res* 1989;484:361–6.
- [92] Theodosis DT, Poulain DA. Maternity leads to morphological synaptic plasticity in the oxytocin system. *Prog Brain Res* 2001;133:49–58.
- [93] Douglas AJ, Johnstone LE, Leng G. Neuroendocrine mechanisms of change in food intake during pregnancy: a potential role for brain oxytocin. *Physiol Behav* 2007;91:352–65.
- [94] Leng G, Caquingueau C, Ludwig M. Priming in Oxytocin Cells and in Gonadotrophs. *Neurochem Res* 2007.
- [95] Ju G, Ma D, Duan XQ. Third ventricular subependymal oxytocin-like immunoreactive neuronal plexus in the rat. *Brain Res Bull* 1992;28:887–96.
- [96] Cloft HJ, Mitchell JA. Immunocytochemical detection of oxytocin in the supraependymal neuronal complex of the golden hamster. *Brain Res* 1994;639:233–9.
- [97] Xiao M, Ding J, Wu L, Han Q, Wang H, Zuo G, Hu G. The distribution of neural nitric oxide synthase-positive cerebrospinal fluid-contacting neurons in the third ventricular wall of male rats and coexistence with vasopressin or oxytocin. *Brain Res* 2005;1038:150–62.
- [98] Robinson AG, Zimmerman EA. Cerebrospinal fluid and ependymal neurophysin. *J Clin Invest* 1973;52:1260–7.
- [99] Vigh B, Manzano e Silva MJ, Frank CL, Vincze C, Czirok SJ, Szabo A, Lukats A, Szel A. The system of cerebrospinal fluid-contacting neurons. Its supposed role in the nonsynaptic signal transmission of the brain. *Histol Histopathol* 2004;19:607–28.
- [100] Vigh B, Vigh-Teichmann I. Actual problems of the cerebrospinal fluid-contacting neurons. *Microsc Res Tech* 1998;41:57–83.
- [101] Cupedo RN, De Weerd H. Serotonergic intraventricular axons in the habenular region; phagocytosis after induced degeneration. *Anat Embryol* 1980;158:213–26.
- [102] Martinez Martinez P, de Weerd H. Supraependymal cell clusters in the rat hypothalamus. *Anat Embryol* 1983;168:371–93.
- [103] Buma P, Nieuwenhuys R. Ultrastructural demonstration of oxytocin and vasopressin release sites in the neural lobe and median eminence of the rat by tannic acid and immunogold methods. *Neurosci Lett* 1987;74:151–7.
- [104] Nicholson C, Sykova E. Extracellular space structure revealed by diffusion analysis. *Trends Neurosci* 1998;21:207–15.
- [105] Sykova E. Extrasynaptic volume transmission and diffusion parameters of the extracellular space. *Neuroscience* 2004;129:861–76.
- [106] Sykova E, Vargova L. Extrasynaptic transmission and the diffusion parameters of the extracellular space. *Neurochem Int* 2008;52:5–13.
- [107] Higuchi T, Tadokoro Y, Honda K, Negoro H. Detailed analysis of blood oxytocin levels during suckling and parturition in the rat. *J Endocrinol* 1986;110:251–6.
- [108] Neumann I, Landgraf R, Bauce L, Pittman QJ. Osmotic responsiveness and cross talk involving oxytocin, but not vasopressin or amino acids, between the supraoptic nuclei in virgin and lactating rats. *J Neurosci* 1995;15:3408–17.
- [109] Moos F, Marganec A, Fontanaud P, Guillou-Duvoix A, Alonso G. Synchronization of oxytocin neurons in suckled rats: possible role of bilateral innervation of hypothalamic supraoptic nuclei by single medullary neurons. *Eur J Neurosci* 2004;20:66–78.
- [110] Neumann ID. Stimuli and consequences of dendritic release of oxytocin within the brain. *Biochem Soc Trans* 2007;35:1252–7.
- [111] Belin V, Moos F. Paired recordings from supraoptic and paraventricular oxytocin cells in suckled rats: recruitment and synchronization. *J Physiol* 1986;377:369–90.
- [112] Rossoni E, Feng J, Tirozzi B, Brown D, Leng G, Moos F. Emergent synchronous bursting of oxytocin neuronal network. *PLoS Comput Biol* 2008;4:e1000123.
- [113] Ebner K, Bosch OJ, Kromer SA, Singewald N, Neumann ID. Release of oxytocin in the rat central amygdala modulates stress-coping behavior and the release of excitatory amino acids. *Neuropsychopharmacology* 2005;30:223–30.
- [114] Larrazolo-Lopez A, Kendrick KM, Aburto-Arciniega M, Arriaga-Avila V, Morimoto S, Frias M, Guevara-Guzman R. Vaginal stimulation enhances social recognition memory in rats via oxytocin release in the olfactory bulb. *Neuroscience* 2008;152:585–93.
- [115] Hughes AM, Everitt BJ, Lightman SL, Todd K. Oxytocin in the central nervous system and sexual behaviour in male rats. *Brain Res* 1987;414:133–7.
- [116] Landgraf R, Neumann I, Schwarzberg H. Central and peripheral release of vasopressin and oxytocin in the conscious rat after osmotic stimulation. *Brain Res* 1988;457:219–25.
- [117] Martinez-Lorenzana G, Espinosa-Lopez L, Carranza M, Aramburo C, Paz-Tres C, Rojas-Piloni G, Condes-Lara M. PVN electrical stimulation prolongs withdrawal latencies and releases oxytocin in cerebrospinal fluid, plasma, and spinal cord tissue in intact and neuropathic rats. *Pain* 2008;140:265–73.
- [118] Morris M, Barnard Jr RR, Sain LE. Osmotic mechanisms regulating cerebrospinal fluid vasopressin and oxytocin in the conscious rat. *Neuroendocrinology* 1984;39:377–83.
- [119] Kendrick KM, Keverne EB, Baldwin BA. Intracerebroventricular oxytocin stimulates maternal behaviour in the sheep. *Neuroendocrinology* 1987;46:56–61.
- [120] Condes-Lara M, Martinez-Lorenzana G, Rodriguez-Jimenez J, Rojas-Piloni G. Paraventricular hypothalamic nucleus stimulation modulates nociceptive responses in dorsal horn wide dynamic range neurons. *Neurosci Lett* 2008;444:199–202.
- [121] Nordmann JJ, Morris JF. Method for quantitating the molecular content of a subcellular organelle: hormone and neurophysin content of newly formed and aged neurosecretory granules. *Proc Natl Acad Sci U S A* 1984;81:180–4.
- [122] Yang J, Yang Y, Chen JM, Liu WY, Wang CH, Lin BC. Effect of oxytocin on acupuncture analgesia in the rat. *Neuropeptides* 2007;41:285–92.
- [123] Freund-Mercier MJ, Stoeckel ME, Palacios JM, Pazos A, Reichhart JM, Porte A, Richard P. Pharmacological characteristics and anatomical distribution of [3H] oxytocin-binding sites in the Wistar rat brain studied by autoradiography. *Neuroscience* 1987;20:599–614.
- [124] Gould BR, Zingg HH. Mapping oxytocin receptor gene expression in the mouse brain and mammary gland using an oxytocin receptor-LacZ reporter mouse. *Neuroscience* 2003;122:155–67.
- [125] Rabhi M, Stoeckel ME, Calas A, Freund-Mercier MJ. Autoradiographic localisation of oxytocin and vasopressin binding sites in the central nervous system of the merione (*Meriones shawi*). *Brain Res Bull* 1999;48:147–63.
- [126] Schorscher-Petcu A, Dupre A, Tribollet E. Distribution of vasopressin and oxytocin binding sites in the brain and upper spinal cord of the common marmoset. *Neurosci Lett* 2009;461:217–22.
- [127] Tribollet E, Barberis C, Dubois-Dauphin M, Dreifuss JJ. Localization and characterization of binding sites for vasopressin and oxytocin in the brain of the guinea pig. *Brain Res* 1992;589:15–23.
- [128] Tribollet E, Charpak S, Schmidt A, Dubois-Dauphin M, Dreifuss JJ. Appearance and transient expression of oxytocin receptors in fetal, infant, and periparturient rat

- brain studied by autoradiography and electrophysiology. *J Neurosci* 1989;9:1764–73.
- [129] Tribollet E, Dubois-Dauphin M, Dreifuss JJ, Barberis C, Jard S. Oxytocin receptors in the central nervous system. Distribution, development, and species differences. *Ann N Y Acad Sci* 1992;652:29–38.
- [130] Vaccari C, Lolait SJ, Ostrowski NL. Comparative distribution of vasopressin V1b and oxytocin receptor messenger ribonucleic acids in brain. *Endocrinology* 1998;139:5015–33.
- [131] Yoshida M, Takayanagi Y, Inoue K, Kimura T, Young LJ, Onaka T, Nishimori K. Evidence that oxytocin exerts anxiolytic effects via oxytocin receptor expressed in serotonergic neurons in mice. *J Neurosci* 2009;29:2259–71.
- [132] Loup F, Tribollet E, Dubois-Dauphin M, Dreifuss JJ. Localization of high-affinity binding sites for oxytocin and vasopressin in the human brain. An autoradiographic study. *Brain Res* 1991;555:220–32.
- [133] Loup F, Tribollet E, Dubois-Dauphin M, Pizzolato G, Dreifuss JJ. Localization of oxytocin binding sites in the human brainstem and upper spinal cord: an autoradiographic study. *Brain Res* 1989;500:223–30.
- [134] Gimpl G, Fahrenholz F. The oxytocin receptor system: structure, function, and regulation. *Physiol Rev* 2001;81:629–83.
- [135] Ostrowski NL. Oxytocin receptor mRNA expression in rat brain: implications for behavioral integration and reproductive success. *Psychoneuroendocrinology* 1998;23:989–1004.
- [136] Lim MM, Wang Z, Olazabal DE, Ren X, Terwilliger EF, Young LJ. Enhanced partner preference in a promiscuous species by manipulating the expression of a single gene. *Nature* 2004;429:754–7.
- [137] Young LJ, Nilsen R, Waymire KG, MacGregor GR, Insel TR. Increased affiliative response to vasopressin in mice expressing the V1a receptor from a monogamous vole. *Nature* 1999;400:766–8.
- [138] Hammock EA, Young LJ. Microsatellite instability generates diversity in brain and sociobehavioral traits. *Science* 2005;308:1630–4.
- [139] Ophir AG, Wolff JO, Phelps SM. Variation in neural V1aR predicts sexual fidelity and space use among male prairie voles in semi-natural settings. *Proc Natl Acad Sci U S A* 2008;105:1249–54.
- [140] Donaldson ZR, Young LJ. Oxytocin, vasopressin, and the neurogenetics of sociality. *Science* 2008;322:900–4.
- [141] Bales KL, Plotsky PM, Young LJ, Lim MM, Grotte N, Ferrer E, Carter CS. Neonatal oxytocin manipulations have long-lasting, sexually dimorphic effects on vasopressin receptors. *Neuroscience* 2007;144:38–45.
- [142] Yoshimura R, Kimura T, Watanabe D, Kiyama H. Differential expression of oxytocin receptor mRNA in the developing rat brain. *Neurosci Res* 1996;24:291–304.
- [143] Insel TR, Gelhard R, Shapiro LE. The comparative distribution of forebrain receptors for neurohypophysial peptides in monogamous and polygamous mice. *Neuroscience* 1991;43:623–30.
- [144] Gimpl G, Burger K, Politowska E, Ciarkowski J, Fahrenholz F. Oxytocin receptors and cholesterol: interaction and regulation. *Exp Physiol* 2000;85:415–9S Spec No.
- [145] Gimpl G, Fahrenholz F. Cholesterol as stabilizer of the oxytocin receptor. *Biochim Biophys Acta* 2002;1564:384–92.
- [146] Thomas A, Shughrue PJ, Merchenthaler I, Amico JA. The effects of progesterone on oxytocin mRNA levels in the paraventricular nucleus of the female rat can be altered by the administration of diazepam or RU486. *J Neuroendocrinol* 1999;11:137–44.
- [147] Windle RJ, Gamble LE, Kershaw YM, Wood SA, Lightman SL, Ingram CD. Gonadal steroid modulation of stress-induced hypothalamo-pituitary-adrenal activity and anxiety behavior: role of central oxytocin. *Endocrinology* 2006;147:2423–31.
- [148] Tribollet E, Audigier S, Dubois-Dauphin M, Dreifuss JJ. Gonadal steroids regulate oxytocin receptors but not vasopressin receptors in the brain of male and female rats. An autoradiographical study. *Brain Res* 1990;511:129–40.
- [149] Patchev VK, Schlosser SF, Hassan AH, Almeida OF. Oxytocin binding sites in rat limbic and hypothalamic structures: site-specific modulation by adrenal and gonadal steroids. *Neuroscience* 1993;57:537–43.
- [150] Liberzon I, Young EA. Effects of stress and glucocorticoids on CNS oxytocin receptor binding. *Psychoneuroendocrinology* 1997;22:411–22.
- [151] Shelat SG, Fluharty SJ, Flanagan-Cato LM. Adrenal steroid regulation of central angiotensin II receptor subtypes and oxytocin receptors in rat brain. *Brain Res* 1998;807:135–46.
- [152] Bale TL, Dorsa DM, Johnston CA. Oxytocin receptor mRNA expression in the ventromedial hypothalamus during the estrous cycle. *J Neurosci* 1995;15:5058–64.
- [153] Caldwell JD, Walker CH, Pedersen CA, Barakat AS, Mason GA. Estrogen increases affinity of oxytocin receptors in the medial preoptic area-anterior hypothalamus. *Peptides* 1994;15:1079–84.
- [154] Agmo A, Choleris E, Kavaliers M, Pfaff DW, Ogawa S. Social and sexual incentive properties of estrogen receptor alpha, estrogen receptor beta, or oxytocin knockout mice. *Genes Brain Behav* 2008;7:70–7.
- [155] Chung SK, McCabe JT, Pfaff DW. Estrogen influences on oxytocin mRNA expression in preoptic and anterior hypothalamic regions studied by *in situ* hybridization. *J Comp Neurol* 1991;307:281–95.
- [156] Nomura M, McKenna E, Korach KS, Pfaff DW, Ogawa S. Estrogen receptor-beta regulates transcript levels for oxytocin and arginine vasopressin in the hypothalamic paraventricular nucleus of male mice. *Brain Res Mol Brain Res* 2002;109:84–94.
- [157] Pfaff DW, Haldar J, Chung SK. *In situ* hybridization for showing hormone effects on oxytocin mRNA in specific populations of hypothalamic neurons and their possible participation in multiplicative hormonal responses. *Ann N Y Acad Sci* 1992;652:347–56.
- [158] Quinones-Jenab V, Jenab S, Ogawa S, Adan RA, Burbach JP, Pfaff DW. Effects of estrogen on oxytocin receptor messenger ribonucleic acid expression in the uterus, pituitary, and forebrain of the female rat. *Neuroendocrinology* 1997;65:9–17.
- [159] Ludwig M, Brown CH, Russell JA, Leng G. Local opioid inhibition and morphine dependence of supraoptic nucleus oxytocin neurones in the rat *in vivo*. *J Physiol* 1997;505(Pt 1):145–52.
- [160] Russell JA, Leng G, Bicknell RJ. Opioid tolerance and dependence in the magnocellular oxytocin system: a physiological mechanism? *Exp Physiol* 1995;80:307–40.
- [161] Proescholdt MG, Hutto B, Brady LS, Herkenham M. Studies of cerebrospinal fluid flow and penetration into brain following lateral ventricle and cisterna magna injections of the tracer [¹⁴C]inulin in rat. *Neuroscience* 2000;95:577–92.
- [162] Herkenham M. Mismatches between neurotransmitter and receptor localizations in brain: observations and implications. *Neuroscience* 1987;23:1–38.
- [163] Dubois-Dauphin M, Tribollet E, Dreifuss JJ. Distribution of neurohypophysial peptides in the guinea pig brain. II. An immunocytochemical study of oxytocin. *Brain Res* 1989;496:66–81.
- [164] Kremarik P, Freund-Mercier MJ, Stoeckel ME. Oxytocin and vasopressin binding sites in the hypothalamus of the rat: histoautoradiographic detection. *Brain Res Bull* 1995;36:195–203.
- [165] Carter CS. Neuroendocrine perspectives on social attachment and love. *Psychoneuroendocrinology* 1998;23:779–818.
- [166] Numan M. Hypothalamic neural circuits regulating maternal responsiveness toward infants. *Behav Cogn Neurosci Rev* 2006;5:163–90.
- [167] Lim MM, Young LJ. Neuropeptide regulation of affiliative behavior and social bonding in animals. *Horm Behav* 2006;50:506–17.
- [168] Neumann ID. Brain mechanisms underlying emotional alterations in the peripartum period in rats. *Depress Anxiety* 2003;17:111–21.
- [169] Insel TR, Young L, Wang Z. Molecular aspects of monogamy. *Ann N Y Acad Sci* 1997;807:302–16.
- [170] McCarthy MM, Kow LM, Pfaff DW. Speculations concerning the physiological significance of central oxytocin in maternal behavior. *Ann N Y Acad Sci* 1992;652:70–82.
- [171] Pedersen CA, Caldwell JD, Peterson G, Walker CH, Mason GA. Oxytocin activation of maternal behavior in the rat. *Ann N Y Acad Sci* 1992;652:58–69.
- [172] Insel TR. Oxytocin—a neuropeptide for affiliation: evidence from behavioral, receptor autoradiographic, and comparative studies. *Psychoneuroendocrinology* 1992;17:3–35.
- [173] Siegel HI. Hormonal basis of maternal behavior in the rat. *Ann N Y Acad Sci* 1986;474:202–15.
- [174] Numan M, Corodimas KP. The effects of paraventricular hypothalamic lesions on maternal behavior in rats. *Physiol Behav* 1985;35:417–25.
- [175] Pedersen CA, Ascher JA, Monroe YL, Prange Jr AJ. Oxytocin induces maternal behavior in virgin female rats. *Science* 1982;216:648–50.
- [176] Wallner B, Dittami J, Machatschke I. Social stimuli cause changes of plasma oxytocin and behavior in guinea pigs. *Biol Res* 2006;39:251–8.
- [177] Numan M. Motivational systems and the neural circuitry of maternal behavior in the rat. *Dev Psychobiol* 2007;49:12–21.
- [178] Poindron P, Levy F, Keller M. Maternal responsiveness and maternal selectivity in domestic sheep and goats: the two facets of maternal attachment. *Dev Psychobiol* 2007;49:54–70.
- [179] Kavaliers M, Choleris E, Agmo A, Braun WJ, Colwell DD, Muglia LJ, Ogawa S, Pfaff DW. Inadvertent social information and the avoidance of parasitized male mice: a role for oxytocin. *Proc Natl Acad Sci U S A* 2006;103:4293–8.
- [180] Bielsky IF, Young LJ. Oxytocin, vasopressin, and social recognition in mammals. *Peptides* 2004;25:1565–74.
- [181] McEwen, B. B. Expansion of olfactory-based social recognition memory research: the roles of vasopressin and oxytocin in social recognition memory. *Adv Pharmacol*. 2004, 50:475–529, 655–708.
- [182] Winslow JT, Insel TR. The social deficits of the oxytocin knockout mouse. *Neuropeptides* 2002;36:221–9.
- [183] Ferguson JN, Aldag JM, Insel TR, Young LJ. Oxytocin in the medial amygdala is essential for social recognition in the mouse. *J Neurosci* 2001;21:8278–85.
- [184] Levy F, Kendrick KM, Goode JA, Guevara-Guzman R, Keverne EB. Oxytocin and vasopressin release in the olfactory bulb of parturient ewes: changes with maternal experience and effects on acetylcholine, gamma-aminobutyric acid, glutamate and noradrenaline release. *Brain Res* 1995;669:197–206.
- [185] Ross HE, Young LJ. Oxytocin and the neural mechanisms regulating social cognition and affiliative behavior. *Front Neuroendocrinol* 2009;30:534–47.
- [186] Ross HE, Freeman SM, Spiegel LL, Ren X, Terwilliger EF, Young LJ. Variation in oxytocin receptor density in the nucleus accumbens has differential effects on affiliative behaviors in monogamous and polygamous voles. *J Neurosci* 2009;29:1312–8.
- [187] Hollander E, Bartz J, Chaplin W, Phillips A, Sumner J, Soorya L, Anagnostou E, Wasserman S. Oxytocin increases retention of social cognition in autism. *Biol Psychiatry* 2007;61:498–503.
- [188] Lim MM, Bielsky IF, Young LJ. Neuropeptides and the social brain: potential rodent models of autism. *Int J Dev Neurosci* 2005;23:235–43.
- [189] Waterhouse L, Fein D, Modahl C. Neurofunctional mechanisms in autism. *Psychol Rev* 1996;103:457–89.
- [190] Young LJ, Pitkow LJ, Ferguson JN. Neuropeptides and social behavior: animal models relevant to autism. *Mol Psychiatry* 2002;7(Suppl 2):S38–9.
- [191] Uvnas-Moberg K. Oxytocin may mediate the benefits of positive social interaction and emotions. *Psychoneuroendocrinology* 1998;23:819–35.

- [192] Franco-Bourland RE. Vasopressinergic, oxytocinergic, and somatostatinergic neuronal activity after adrenalectomy and immobilization stress. *Neurochem Res* 1998;23:695–701.
- [193] Grassi M, Drago F. Effects of oxytocin on emotional stress and stress-induced gastric lesions. *J Physiol Paris* 1993;87:261–4.
- [194] Muir JL, Pfister HP. Psychological stress and oxytocin treatment during pregnancy affect central norepinephrine, dopamine and serotonin in lactating rats. *Int J Neurosci* 1989;48:191–203.
- [195] Neumann ID. Alterations in behavioral and neuroendocrine stress coping strategies in pregnant, parturient and lactating rats. *Prog Brain Res* 2001;133:143–52.
- [196] Zou CJ, Onaka T, Yagi K. Role of adrenoceptors in vasopressin, oxytocin and prolactin responses to conditioned fear stimuli in the rat. *J Neuroendocrinol* 1998;10:905–10.
- [197] Agren G, Lundeberg T. Energy conservation in stressed rats exposed to an oxytocin-injected cage mate. *NeuroReport* 2002;13:1453–7.
- [198] Agren G, Olsson C, Uvnas-Moberg K, Lundeberg T. Olfactory cues from an oxytocin-injected male rat can reduce energy loss in its cagemates. *NeuroReport* 1997;8:2551–5.
- [199] Born J, Lange T, Kern W, McGregor GP, Bickel U, Fehm HL. Sniffing neuropeptides: a transnasal approach to the human brain. *Nat Neurosci* 2002;5:514–6.
- [200] Charlton ST, Whetstone J, Fayinka ST, Read KD, Illum L, Davis SS. Evaluation of direct transport pathways of glycine receptor antagonists and an angiotensin antagonist from the nasal cavity to the central nervous system in the rat model. *Pharm Res* 2008;25:1531–43.
- [201] Dhuria SV, Hanson LR, Frey 2nd WH. Intranasal delivery to the central nervous system: mechanisms and experimental considerations. *J Pharm Sci* 2010;99:1654–73.
- [202] Dhuria SV, Hanson LR, Frey 2nd WH. Intranasal drug targeting of hypocretin-1 (orexin-A) to the central nervous system. *J Pharm Sci* 2009;98:2501–15.
- [203] Illum L. The nasal delivery of peptides and proteins. *Trends Biotechnol* 1991;9:284–9.
- [204] Illum L. Transport of drugs from the nasal cavity to the central nervous system. *Eur J Pharm Sci* 2000;11:1–18.
- [205] Thorne RG, Emory CR, Ala TA, Frey 2nd WH. Quantitative analysis of the olfactory pathway for drug delivery to the brain. *Brain Res* 1995;692:278–82.
- [206] Thorne RG, Frey 2nd WH. Delivery of neurotrophic factors to the central nervous system: pharmacokinetic considerations. *Clin Pharmacokinet* 2001;40:907–46.
- [207] Thorne RG, Pronk GJ, Padmanabhan V, Frey 2nd WH. Delivery of insulin-like growth factor-I to the rat brain and spinal cord along olfactory and trigeminal pathways following intranasal administration. *Neuroscience* 2004;127:481–96.
- [208] McGregor IS, Callaghan PD, Hunt GE. From ultrasocial to antisocial: a role for oxytocin in the acute reinforcing effects and long-term adverse consequences of drug use? *Br J Pharmacol* 2008;154:358–68.
- [209] Macdonald K, Macdonald TM. The peptide that binds: a systematic review of oxytocin and its prosocial effects in humans. *Harv Rev Psychiatry* 2010;18:1–21.
- [210] Fischer-Shofty M, Shamay-Tsoory SG, Harari H, Levkovitz Y. The effect of intranasal administration of oxytocin on fear recognition. *Neuropsychologia* 2010;48:179–84.
- [211] Guastella AJ, Howard AL, Dadds MR, Mitchell P, Carson DS. A randomized controlled trial of intranasal oxytocin as an adjunct to exposure therapy for social anxiety disorder. *Psychoneuroendocrinology* 2009;34:917–23.
- [212] Di Simplicio M, Massey-Chase R, Cowen PJ, Harmer CJ. Oxytocin enhances processing of positive versus negative emotional information in healthy male volunteers. *J Psychopharmacol* 2009;23:241–8.
- [213] Heim C, Young LJ, Newport DJ, Mletzko T, Miller AH, Nemeroff CB. Lower CSF oxytocin concentrations in women with a history of childhood abuse. *Mol Psychiatry* 2009;14:954–8.
- [214] Kosfeld M, Heinrichs M, Zak PJ, Fischbacher U, Fehr E. Oxytocin increases trust in humans. *Nature* 2005;435:673–6.
- [215] Domes G, Heinrichs M, Michel A, Berger C, Herpertz SC. Oxytocin improves "mind-reading" in humans. *Biol Psychiatry* 2007;61:731–3.
- [216] Illum L. Nasal drug delivery—possibilities, problems and solutions. *J Control Release* 2003;87:187–98.
- [217] Illum L. Is nose-to-brain transport of drugs in man a reality? *J Pharm Pharmacol* 2004;56:3–17.
- [218] Argiolas A, Melis MR, Gessa GL. Oxytocin: an extremely potent inducer of penile erection and yawning in male rats. *Eur J Pharmacol* 1986;130:265–72.
- [219] Melis MR, Argiolas A, Gessa GL. Evidence that apomorphine induces penile erection and yawning by releasing oxytocin in the central nervous system. *Eur J Pharmacol* 1989;164:565–70.
- [220] Tang Y, Rampin O, Calas A, Facchinetti P, Giuliano F. Oxytocinergic and serotonergic innervation of identified lumbosacral nuclei controlling penile erection in the male rat. *Neuroscience* 1998;82:241–54.
- [221] Veronneau-Longueville F, Rampin O, Freund-Mercier MJ, Tang Y, Calas A, Marson L, McKenna KE, Stoeckel ME, Benoit G, Giuliano F. Oxytocinergic innervation of autonomic nuclei controlling penile erection in the rat. *Neuroscience* 1999;93:1437–47.
- [222] Giuliano F, Bernabe J, McKenna K, Longueville F, Rampin O. Spinal proerectile effect of oxytocin in anesthetized rats. *Am J Physiol Regul Integr Comp Physiol* 2001;280:R1870–7.
- [223] Argiolas A, Melis MR. Central control of penile erection: role of the paraventricular nucleus of the hypothalamus. *Prog Neurobiol* 2005;76:1–21.
- [224] Succu S, Sanna F, Melis T, Boi A, Argiolas A, Melis MR. Stimulation of dopamine receptors in the paraventricular nucleus of the hypothalamus of male rats induces penile erection and increases extra-cellular dopamine in the nucleus accumbens: involvement of central oxytocin. *Neuropharmacology* 2007;52:1034–43.
- [225] Zhang XH, Filippi S, Vignozzi L, Morelli A, Mancina R, Luconi M, Donati S, Marini M, Vannelli GB, Forti G, Maggi M. Identification, localization and functional in vitro and in vivo activity of oxytocin receptor in the rat penis. *J Endocrinol* 2005;184:567–76.
- [226] Argiolas A. Oxytocin stimulation of penile erection. Pharmacology, site, and mechanism of action. *Ann N Y Acad Sci* 1992;652:194–203.
- [227] Melis MR, Succu S, Cocco C, Caboni E, Sanna F, Boi A, Ferri GL, Argiolas A. Oxytocin induces penile erection when injected into the ventral subiculum: role of nitric oxide and glutamic acid. *Neuropharmacology* 2010;58:1153–60.
- [228] Melis MR, Succu S, Sanna F, Boi A, Argiolas A. Oxytocin injected into the ventral subiculum or the posteromedial cortical nucleus of the amygdala induces penile erection and increases extracellular dopamine levels in the nucleus accumbens of male rats. *Eur J Neurosci* 2009;30:1349–57.
- [229] Arletti R, Benelli A, Bertolini A. Sexual behavior of aging male rats is stimulated by oxytocin. *Eur J Pharmacol* 1990;179:377–81.
- [230] Arletti R, Benelli A, Bertolini A. Oxytocin involvement in male and female sexual behavior. *Ann N Y Acad Sci* 1992;652:180–93.
- [231] Arletti R, Bazzani C, Castelli M, Bertolini A. Oxytocin improves male copulatory performance in rats. *Horm Behav* 1985;19:14–20.
- [232] Ishak WW, Berman DS, Peters A. Male anorgasmia treated with oxytocin. *J Sex Med* 2008;5:1022–4.
- [233] Filippi S, Vignozzi L, Vannelli GB, Ledda F, Forti G, Maggi M. Role of oxytocin in the ejaculatory process. *J Endocrinol Invest* 2003;26:82–6.
- [234] Kunz G, Beil D, Huppert P, Leyendecker G. Oxytocin—a stimulator of directed sperm transport in humans. *Reprod Biomed Online* 2007;14:32–9.
- [235] Zervomanolakis I, Ott HW, Hadziomerovic D, Mattle V, Seeber BE, Virgolini I, Heute D, Kissler S, Leyendecker G, Wildt L. Physiology of upward-transport in the human female genital tract. *Ann N Y Acad Sci* 2007.
- [236] Moody KM, Steinman JL, Komisaruk BR, Adler NT. Pelvic neurectomy blocks oxytocin-facilitated sexual receptivity in rats. *Physiol Behav* 1994;56:1057–60.
- [237] Arletti R, Bertolini A. Oxytocin acts as an antidepressant in two animal models of depression. *Life Sci* 1987;41:1725–30.
- [238] Olivier B, Chan JS, Pattij T, de Jong TR, Oosting RS, Veening JG, Waldinger MD. Psychopharmacology of male rat sexual behavior: modeling human sexual dysfunctions? *Int J Impot Res* 2006;18(Suppl 1):S14–23.
- [239] Cantor JM, Binik YM, Pfaus JG. Chronic fluoxetine inhibits sexual behavior in the male rat: reversal with oxytocin. *Psychopharmacology (Berl)* 1999;144:355–62.
- [240] de Jong TR, Veening JG, Olivier B, Waldinger MD. Oxytocin involvement in SSRI-induced delayed ejaculation: a review of animal studies. *J Sex Med* 2007;4:14–28.
- [241] Argiolas A, Gessa GL. Central functions of oxytocin. *Neurosci Biobehav Rev* 1991;15:217–31.
- [242] Stoneham MD, Everitt BJ, Hansen S, Lightman SL, Todd K. Oxytocin and sexual behaviour in the male rat and rabbit. *J Endocrinol* 1985;107:97–106.
- [243] Mahalati K, Okanoya K, Witt DM, Carter CS. Oxytocin inhibits male sexual behavior in prairie voles. *Pharmacol Biochem Behav* 1991;39:219–22.
- [244] Carter CS. Oxytocin and sexual behavior. *Neurosci Biobehav Rev* 1992;16:131–44.
- [245] Witt DM, Carter CS, Walton DM. Central and peripheral effects of oxytocin administration in prairie voles (*Microtus ochrogaster*). *Pharmacol Biochem Behav* 1990;15:217–31.
- [246] Argiolas A, Melis MR. The role of oxytocin and the paraventricular nucleus in the sexual behaviour of male mammals. *Physiol Behav* 2004;83:309–17.
- [247] Chen K, Chang LS. Oxytocinergic neurotransmission at the hippocampus in the central neural regulation of penile erection in the rat. *Urology* 2001;58:107–12.
- [248] Pedersen CA, Boccia ML. Vasopressin interactions with oxytocin in the control of female sexual behavior. *Neuroscience* 2006;139:843–51.
- [249] Pedersen CA, Boccia ML. Oxytocin maintains as well as initiates female sexual behavior: effects of a highly selective oxytocin antagonist. *Horm Behav* 2002;41:170–7.
- [250] Gorzalka BB, Lester GL. Oxytocin-induced facilitation of lordosis behaviour in rats is progesterone-dependent. *Neuropeptides* 1987;10:55–65.
- [251] Caldwell JD, Prange Jr AJ, Pedersen CA. Oxytocin facilitates the sexual receptivity of estrogen-treated female rats. *Neuropeptides* 1986;7:175–89.
- [252] Arletti R, Bertolini A. Oxytocin stimulates lordosis behavior in female rats. *Neuropeptides* 1985;6:247–53.
- [253] Caldwell JD. Central oxytocin and female sexual behavior. *Ann N Y Acad Sci* 1992;652:166–79.
- [254] Gelez H, Poirier S, Facchinetti P, Allers KA, Wayman C, Alexandre L, Giuliano F. Neuroanatomical Evidence for a Role of Central Melanocortin-4 Receptors and Oxytocin in the Efferent Control of the Rodent Clitoris and Vagina. *J Sex Med* 2010.
- [255] Caldwell JD, Jirikowski GF, Greer ER, Stumpf WE, Pedersen CA. Ovarian steroids and sexual interaction alter oxytocinergic content and distribution in the basal forebrain. *Brain Res* 1988;446:236–44.
- [256] Flanagan LM, Pfaus JG, Pfaff DW, McEwen BS. Induction of FOS immunoreactivity in oxytocin neurons after sexual activity in female rats. *Neuroendocrinology* 1993;58:352–8.
- [257] McCarthy MM, Kleopoulos SP, Mobbs CV, Pfaff DW. Infusion of antisense oligodeoxynucleotides to the oxytocin receptor in the ventromedial hypothalamus reduces estrogen-induced sexual receptivity and oxytocin receptor binding in the female rat. *Neuroendocrinology* 1994;59:432–40.
- [258] Mena F, Aguayo D, Pacheco P, Morales MT. Central effects of catecholamines upon mammary contractility in rats are neurally mediated. *Neuroendocrinology* 1995;61:722–30.

- [259] Fenelon VS, Poulain DA, Theodosis DT. Oxytocin neuron activation and Fos expression: a quantitative immunocytochemical analysis of the effect of lactation, parturition, osmotic and cardiovascular stimulation. *Neuroscience* 1993;53:77–89.
- [260] Terenzi MG, Ingram CD. Oxytocin-induced excitation of neurones in the rat central and medial amygdaloid nuclei. *Neuroscience* 2005;134:345–54.
- [261] Lambert RC, Moos FC, Richard P. Action of endogenous oxytocin within the paraventricular or supraoptic nuclei: a powerful link in the regulation of the bursting pattern of oxytocin neurons during the milk-ejection reflex in rats. *Neuroscience* 1993;57:1027–38.
- [262] Young LJ, Muns S, Wang Z, Insel TR. Changes in oxytocin receptor mRNA in rat brain during pregnancy and the effects of estrogen and interleukin-6. *J Neuroendocrinol* 1997;9:859–65.
- [263] Neumann I, Russell JA, Landgraf R. Oxytocin and vasopressin release within the supraoptic and paraventricular nuclei of pregnant, parturient and lactating rats: a microdialysis study. *Neuroscience* 1993;53:65–75.
- [264] Tyzio R, Cossart R, Khalilov I, Minlebaev M, Hubner CA, Represa A, Ben-Ari Y, Khazipov R. Maternal oxytocin triggers a transient inhibitory switch in GABA signaling in the fetal brain during delivery. *Science* 2006;314:1788–92.
- [265] Neumann ID, Torner L, Wigger A. Brain oxytocin: differential inhibition of neuroendocrine stress responses and anxiety-related behaviour in virgin, pregnant and lactating rats. *Neuroscience* 2000;95:567–75.
- [266] Febo M, Shields J, Ferris CF, King JA. Oxytocin modulates unconditioned fear response in lactating dams: an fMRI study. *Brain Res* 2009;1302:183–93.
- [267] Da Costa AP, Guevara-Guzman RG, Ohkura S, Goode JA, Kendrick KM. The role of oxytocin release in the paraventricular nucleus in the control of maternal behaviour in the sheep. *J Neuroendocrinol* 1996;8:163–77.
- [268] McCann MJ, Rogers RC. Oxytocin excites gastric-related neurones in rat dorsal vagal complex. *J Physiol* 1990;428:95–108.
- [269] Cruz MT, Murphy EC, Sahibzada N, Verbalis JG, Gillis RA. A reevaluation of the effects of stimulation of the dorsal motor nucleus of the vagus on gastric motility in the rat. *Am J Physiol Regul Integr Comp Physiol* 2007;292:R291–307.
- [270] Asad M, Shewade DG, Koumaravelou K, Abraham BK, Vasu S, Ramaswamy S. Gastric antisecretory and antiulcer activity of oxytocin in rats and guinea pigs. *Life Sci* 2001;70:17–24.
- [271] Siaud P, Puech R, Assenmacher I, Alonso G. Microinjection of oxytocin into the dorsal vagal complex decreases pancreatic insulin secretion. *Brain Res* 1991;546:190–4.
- [272] Sclafani A, Rinaman L, Vollmer RR, Amico JA. Oxytocin knockout mice demonstrate enhanced intake of sweet and non-sweet carbohydrate solutions. *Am J Physiol Regul Integr Comp Physiol* 2007.
- [273] Blevins JE, Eakin TJ, Murphy JA, Schwartz MW, Baskin DG. Oxytocin innervation of caudal brainstem nuclei activated by cholecystokinin. *Brain Res* 2003;993:30–41.
- [274] Blevins JE, Schwartz MW, Baskin DG. Evidence that paraventricular nucleus oxytocin neurons link hypothalamic leptin action to caudal brain stem nuclei controlling meal size. *Am J Physiol Regul Integr Comp Physiol* 2004;287:R87–96.
- [275] Sabatier N, Rowe I, Leng C. Central release of oxytocin and the ventromedial hypothalamus. *Biochem Soc Trans* 2007;35:1247–51.
- [276] Meisenberg G, Simmons WH. Behavioral effects of intracerebroventricularly administered neurohypophyseal hormone analogs in mice. *Pharmacol Biochem Behav* 1982;16:819–25.
- [277] Pedersen CA, Caldwell JD, Drago F, Noonan LR, Peterson G, Hood LE, Prange Jr AJ. Grooming behavioral effects of oxytocin. *Pharmacology, ontogeny, and comparisons with other nonapeptides*. *Ann N Y Acad Sci* 1988;525:245–56.
- [278] Van Erp AM, Kruk MR, De Kloet ER. Induction of grooming in resting rats by intracerebroventricular oxytocin but not by adrenocorticotrophic hormone-(1–24) and alpha-melanocyte-stimulating hormone. *Eur J Pharmacol* 1993;232:217–21.
- [279] Van Wimersma Greidanus TB, Kroodsmas JM, Pot ML, Stevens M, Maigret C. Neurohypophyseal hormones and excessive grooming behaviour. *Eur J Pharmacol* 1990;187:1–8.
- [280] Amico JA, Vollmer RR, Karam JR, Lee PR, Li X, Koenig JJ, McCarthy MM. Centrally administered oxytocin elicits exaggerated grooming in oxytocin null mice. *Pharmacol Biochem Behav* 2004;78:333–9.
- [281] Kaltwasser MT, Crawley JN. Oxytocin and cholecystokinin induce grooming behavior in the ventral tegmentum of the rat. *Brain Res* 1987;426:1–7.
- [282] van Erp AM, Kruk MR, Veening JG, Roeling TA, Meelis W. Neuronal substrate of electrically induced grooming in the PVH of the rat: involvement of oxytocinergic systems? *Physiol Behav* 1995;57:881–5.
- [283] Drago F, Sarnyai Z, D'Agata V. The inhibition of oxytocin-induced grooming by a specific receptor antagonist. *Physiol Behav* 1991;50:533–6.
- [284] Arletti R, Benelli A, Bertolini A. Influence of oxytocin on nociception and morphine antinociception. *Neuropeptides* 1993;24:125–9.
- [285] Ge Y, Lundeberg T, Yu LC. Blockade effect of mu and kappa opioid antagonists on the anti-nociception induced by intra-periaqueductal grey injection of oxytocin in rats. *Brain Res* 2002;927:204–7.
- [286] Gu XL, Yu LC. Involvement of opioid receptors in oxytocin-induced antinociception in the nucleus accumbens of rats. *J Pain* 2007;8:85–90.
- [287] Millan MJ, Schmauss C, Millan MH, Herz A. Vasopressin and oxytocin in the rat spinal cord: analysis of their role in the control of nociception. *Brain Res* 1984;309:384–8.
- [288] Yang J, Yang Y, Chen JM, Liu WY, Wang CH, Lin BC. Central oxytocin enhances antinociception in the rat. *Peptides* 2007;28:1113–9.
- [289] Backman SB, Henry JL. Effects of oxytocin and vasopressin on thoracic sympathetic preganglionic neurones in the cat. *Brain Res Bull* 1984;13:679–84.
- [290] Sansone GR, Komisaruk BR. Evidence that oxytocin is an endogenous stimulator of autonomic sympathetic preganglionics: the pupillary dilatation response to vaginocervical stimulation in the rat. *Brain Res* 2001;898:265–71.
- [291] Busija DW, Khreis I, Chen J. Prostanoids promote pial arteriolar dilation and mask constriction to oxytocin in piglets. *Am J Physiol* 1993;264:H1023–7.
- [292] Swanson LW. Cerebral hemisphere regulation of motivated behavior. *Brain Res* 2000;886:113–64.
- [293] Choleris E, Gustafsson JA, Korach KS, Muglia LJ, Pfaff DW, Ogawa S. An estrogen-dependent four-gene micronet regulating social recognition: a study with oxytocin and estrogen receptor-alpha and -beta knockout mice. *Proc Natl Acad Sci U S A* 2003;100:6192–7.
- [294] Dellovade TL, Zhu YS, Pfaff DW. Thyroid hormones and estrogen affect oxytocin gene expression in hypothalamic neurons. *J Neuroendocrinol* 1999;11:1–10.
- [295] Kavaliers M, Agmo A, Choleris E, Gustafsson JA, Korach KS, Muglia LJ, Pfaff DW, Ogawa S. Oxytocin and estrogen receptor alpha and beta knockout mice provide discriminably different odor cues in behavioral assays. *Genes Brain Behav* 2004;3:189–95.
- [296] Kow LM, Johnson AE, Ogawa S, Pfaff DW. Electrophysiological actions of oxytocin on hypothalamic neurons in vitro: neuropharmacological characterization and effects of ovarian steroids. *Neuroendocrinology* 1991;54:526–35.
- [297] Schumacher M, Coirini H, Flanagan LM, Frankfurt M, Pfaff DW, McEwen BS. Ovarian steroid modulation of oxytocin receptor binding in the ventromedial hypothalamus. *Ann N Y Acad Sci* 1992;652:374–86.
- [298] Ragnauth AK, Goodwillie A, Brewer C, Muglia LJ, Pfaff DW, Kow LM. Vasopressin stimulates ventromedial hypothalamic neurons via oxytocin receptors in oxytocin gene knockout male and female mice. *Neuroendocrinology* 2004;80:92–9.
- [299] Schumacher M, Coirini H, Johnson AE, Flanagan LM, Frankfurt M, Pfaff DW, McEwen BS. The oxytocin receptor: a target for steroid hormones. *Regul Pept* 1993;45:115–9.
- [300] Schumacher M, Coirini H, Pfaff DW, McEwen BS. Behavioral effects of progesterone associated with rapid modulation of oxytocin receptors. *Science* 1990;250:691–4.
- [301] Coolen LM, Olivier B, Peters HJ, Veening JG. Demonstration of ejaculation-induced neural activity in the male rat brain using 5-HT1A agonist 8-OH-DPAT. *Physiol Behav* 1997;62:881–91.
- [302] Coolen LM, Peters HJ, Veening JG. Fos immunoreactivity in the rat brain following consummatory elements of sexual behavior: a sex comparison. *Brain Res* 1996;738:67–82.
- [303] Veening JG, Coolen LM. Neural activation following sexual behavior in the male and female rat brain. *Behav Brain Res* 1998;92:181–93.
- [304] Calle M, Claassen IE, Veening JG, Kozic T, Roubos EW, Barendregt HP. Opioid peptides, CRF, and urocortin in cerebrosal fluid-contacting neurons in *Xenopus laevis*. *Ann N Y Acad Sci* 2005;1040:249–52.
- [305] Roeling TA, Kruk MR, Schuurmans R, Veening JG. Behavioural responses of bicuculline methiodide injections into the ventral hypothalamus of freely moving, socially interacting rats. *Brain Res* 1993;615:121–7.
- [306] Roeling TA, Veening JG, Kruk MR, Peters JP, Vermelis ME, Nieuwenhuys R. Efferent connections of the hypothalamic "aggression area" in the rat. *Neuroscience* 1994;59:1001–24.
- [307] Halasz J, Liposits Z, Meelis W, Kruk MR, Haller J. Hypothalamic attack area-mediated activation of the forebrain in aggression. *NeuroReport* 2002;13:1267–70.
- [308] Kruk MR, Van der Laan CE, Mos J, Van der Poel AM, Meelis W, Olivier B. Comparison of aggressive behaviour induced by electrical stimulation in the hypothalamus of male and female rats. *Prog Brain Res* 1984;61:303–14.
- [309] Kruk MR, Van der Poel AM, Meelis W, Hermans J, Mostert PG, Mos J, Lohman AH. Discriminant analysis of the localization of aggression-inducing electrode placements in the hypothalamus of male rats. *Brain Res* 1983;260:61–79.
- [310] Lammers JH, Kruk MR, Meelis W, van der Poel AM. Hypothalamic substrates for brain stimulation-induced attack, teeth-chattering and social grooming in the rat. *Brain Res* 1988;449:311–27.
- [311] Siegel A, Roeling TA, Gregg TR, Kruk MR. Neuropharmacology of brain-stimulation-evoked aggression. *Neurosci Biobehav Rev* 1999;23:359–89.
- [312] Dubois-Dauphin M, Raggenbass M, Widmer H, Tribollet E, Dreifuss JJ. Morphological and electrophysiological evidence for postsynaptic localization of functional oxytocin receptors in the rat dorsal motor nucleus of the vagus nerve. *Brain Res* 1992;575:124–31.
- [313] Komisaruk BR, Sansone G. Neural pathways mediating vaginal function: the vagus nerves and spinal cord oxytocin. *Scand J Psychol* 2003;44:241–50.
- [314] Borison HL, Borison R, McCarthy LE. Brain stem penetration by horseradish peroxidase from the cerebrospinal fluid spaces in the cat. *Exp Neurol* 1980;69:271–89.
- [315] Ferguson IA, Schweitzer JB, Bartlett PF, Johnson Jr EM. Receptor-mediated retrograde transport in CNS neurons after intraventricular administration of NGF and growth factors. *J Comp Neurol* 1991;313:680–92.
- [316] Konsman JP, Tridon V, Dantzer R. Diffusion and action of intracerebroventricularly injected interleukin-1 in the CNS. *Neuroscience* 2000;101:957–67.
- [317] Caraty A, Delaere B, Chesneau D, Fabre-Nys C. Sequential role of e2 and GnRH for the expression of estrous behavior in ewes. *Endocrinology* 2002;143:139–45.
- [318] Caraty A, Skinner DC. Gonadotropin-releasing hormone in third ventricular cerebrospinal fluid: endogenous distribution and exogenous uptake. *Endocrinology* 2008;149:5227–34.
- [319] Bruni JE. Ependymal development, proliferation, and functions: a review. *Microsc Res Tech* 1998;41:2–13.
- [320] Bruni JE, Reddy K. Ependyma of the central canal of the rat spinal cord: a light and transmission electron microscopic study. *J Anat* 1987;152:55–70.

- [321] Bruni JE, Clattenburg RE, Montemurro DG. Ependymal tanycytes of the rabbit third ventricle: a scanning electron microscopic study. *Brain Res* 1974;73:145–50.
- [322] Guerra M, Blazquez JL, Peruzzo B, Pelaez B, Rodriguez S, Toranzo D, Pastor F, Rodriguez EM. Cell organization of the rat pars tuberalis. Evidence for open communication between pars tuberalis cells, cerebrospinal fluid and tanycytes. *Cell Tissue Res* 2009;339:359–81.
- [323] Rodriguez EM, Blazquez JL, Pastor FE, Pelaez B, Pena P, Peruzzo B, Amat P. Hypothalamic tanycytes: a key component of brain-endocrine interaction. *Int Rev Cytol* 2005;247:89–164.
- [324] Peruzzo B, Pastor FE, Blazquez JL, Amat P, Rodriguez EM. Polarized endocytosis and transcytosis in the hypothalamic tanycytes of the rat. *Cell Tissue Res* 2004;317:147–64.
- [325] Zec N, Kinney HC. Anatomic relationships of the human nucleus paragigantocellularis lateralis: a Dil labeling study. *Auton Neurosci* 2001;89:110–24.
- [326] Felten DL, Cummings JP, Burnett BT. Ontogeny of caudal fourth ventricular tanycytes in the rabbit brain: a Golgi study. *Anat Rec* 1981;200:321–30.
- [327] Felten DL, Harrigan P, Burnett BT, Cummings JP. Fourth ventricular tanycytes: a possible relationship with monoaminergic nuclei. *Brain Res Bull* 1981;6:427–36.
- [328] Felten DL, Harrigan P. Dendrite bundles in nuclei raphe dorsalis and centralis superior of the rabbit: a possible substrate for local control of serotonergic neurons. *Neurosci Lett* 1980;16:275–80.
- [329] Cummings JP, Felten DL. A raphe dendrite bundle in the rabbit medulla. *J Comp Neurol* 1979;183:1–23.
- [330] Torterolo P, Lagos P, Sampogna S, Chase MH. Melanin-concentrating hormone (MCH) immunoreactivity in non-neuronal cells within the raphe nuclei and subventricular region of the brainstem of the cat. *Brain Res* 2008;1210:163–78.
- [331] Feng C, Baryshnikova LM, von Bartheld CS. A novel systemic route for specific accumulation of Nerve Growth Factor in the Central Nervous System. Society for Neuroscience; 2007. San Diego, CA, USA.
- [332] Miyan JA, Nabiyouni M, Zenda M. Development of the brain: a vital role for cerebrospinal fluid. *Can J Physiol Pharmacol* 2003;81:317–28.
- [333] De Wied D, Diamant M, Fodor M. Central nervous system effects of the neurohypophysial hormones and related peptides. *Front Neuroendocrinol* 1993;14:251–302.
- [334] Febo M, Numan M, Ferris CF. Functional magnetic resonance imaging shows oxytocin activates brain regions associated with mother-pup bonding during suckling. *J Neurosci* 2005;25:11637–44.
- [335] Lincoln DW, Hentzen K, Hin T, van der Schoot P, Clarke G, Summerlee AJ. Sleep: a prerequisite for reflex milk ejection in the rat. *Exp Brain Res* 1980;38:151–62.
- [336] Silakov VL, Nikitin VS, Moiseeva LA, Losev SS, Perepelkin PD. The comparative action of relanin and oxytocin on higher nervous activity in lower monkeys. *Zh Vyssh Nerv Deiat Im I P Pavlova* 1992;42:734–42.
- [337] Pietrowsky R, Braun D, Fehm HL, Pauschinger P, Born J. Vasopressin and oxytocin do not influence early sensory processing but affect mood and activation in man. *Peptides* 1991;12:1385–91.
- [338] Baumgartner T, Heinrichs M, Vonlanthen A, Fischbacher U, Fehr E. Oxytocin shapes the neural circuitry of trust and trust adaptation in humans. *Neuron* 2008;58:639–50.
- [339] Delgado MR. Fool me once, shame on you; fool me twice, shame on oxytocin. *Neuron* 2008;58:470–1.
- [340] Petrovic P, Kalisch R, Singer T, Dolan RJ. Oxytocin attenuates affective evaluations of conditioned faces and amygdala activity. *J Neurosci* 2008;28:6607–15.
- [341] Allman JM, Watson KK, Tetreault NA, Hakeem AY. Intuition and autism: a possible role for Von Economo neurons. *Trends Cogn Sci* 2005;9:367–73.
- [342] Fajardo C, Escobar MI, Buritica E, Arteaga G, Umbarila J, Casanova MF, Pimienta H. Von Economo neurons are present in the dorsolateral (dysgranular) prefrontal cortex of humans. *Neurosci Lett* 2008;435:215–8.
- [343] Viskontas IV, Possin KL, Miller BL. Symptoms of frontotemporal dementia provide insights into orbitofrontal cortex function and social behavior. *Ann N Y Acad Sci* 2007;1121:528–45.
- [344] Meyer-Lindenberg A. Impact of prosocial neuropeptides on human brain function. *Prog Brain Res* 2008;170:463–70.
- [345] Opar A. Search for potential autism treatments turns to 'trust hormone'. *Nat Med* 2008;14:353.
- [346] Rimmele U, Hediger K, Heinrichs M, Klaver P. Oxytocin makes a face in memory familiar. *J Neurosci* 2009;29:38–42.
- [347] Yu GZ, Kaba H, Okutani F, Takahashi S, Higuchi T, Seto K. The action of oxytocin originating in the hypothalamic paraventricular nucleus on mitral and granule cells in the rat main olfactory bulb. *Neuroscience* 1996;72:1073–82.
- [348] Meddle SL, Bishop VR, Koumassi E, van Leeuwen FW, Douglas AJ. Dynamic changes in oxytocin receptor expression and activation at parturition in the rat brain. *Endocrinology* 2007;148:5095–104.
- [349] Hatterer E, Davoust N, Didier-Bazes M, Vuillat C, Malcus C, Belin MF, Nataf S. How to drain without lymphatics? Dendritic cells migrate from the cerebrospinal fluid to the B-cell follicles of cervical lymph nodes. *Blood* 2006;107:806–12.
- [350] Moffatt CA. Steroid hormone modulation of olfactory processing in the context of socio-sexual behaviors in rodents and humans. *Brain Res Brain Res Rev* 2003;43:192–206.
- [351] Purdon SE, Klein S, Flor-Henry P. Menstrual effects on asymmetrical olfactory acuity. *J Int Neuropsychol Soc* 2001;7:703–9.
- [352] Wesson DW, Keller M, Douhard Q, Baum MJ, Bakker J. Enhanced urinary odor discrimination in female aromatase knockout (ArKO) mice. *Horm Behav* 2006;49:580–6.
- [353] Doane DF, Lawson MA, Meade JR, Kotz CM, Beverly JL. Orexin-induced feeding requires NMDA receptor activation in the perifornical region of the lateral hypothalamus. *Am J Physiol Regul Integr Comp Physiol* 2007;293:R1022–6.
- [354] Edwards CM, Abusnana S, Sunter D, Murphy KG, Ghatei MA, Bloom SR. The effect of the orexins on food intake: comparison with neuropeptide Y, melanin-concentrating hormone and galanin. *J Endocrinol* 1999;160:R7–R12.
- [355] Jang EH, Moon JS, Ko JH, Ahn CW, Lee HH, Shin JK, Park CS, Kang JH. Novel black soy peptides with antiobesity effects: activation of leptin-like signaling and AMP-activated protein kinase. *Int J Obes (Lond)* 2008.
- [356] Samson WK, Murphy TC, Robison D, Vargas T, Tau E, Chang JK. A 35 amino acid fragment of leptin inhibits feeding in the rat. *Endocrinology* 1996;137:5182–5.
- [357] Zhu JN, Guo CL, Li HZ, Wang JJ. Dorsomedial hypothalamic nucleus neurons integrate important peripheral feeding-related signals in rats. *J Neurosci Res* 2007;85:3193–204.
- [358] Aime P, Duchamp-Viret P, Chaput MA, Savigner A, Mahfouz M, Julliard AK. Fasting increases and satiation decreases olfactory detection for a neutral odor in rats. *Behav Brain Res* 2007;179:258–64.
- [359] Julliard AK, Chaput MA, Apelbaum A, Aime P, Mahfouz M, Duchamp-Viret P. Changes in rat olfactory detection performance induced by orexin and leptin mimicking fasting and satiation. *Behav Brain Res* 2007;183:123–9.
- [360] Baly C, Aioun J, Badonnel K, Lacroix MC, Durieux D, Schlegel C, Salesse R, Caillol M. Leptin and its receptors are present in the rat olfactory mucosa and modulated by the nutritional status. *Brain Res* 2007;1129:130–41.
- [361] Levasseur G, Baly C, Grebert D, Durieux D, Salesse R, Caillol M. Anatomical and functional evidence for a role of arginine-vasopressin (AVP) in rat olfactory epithelium cells. *Eur J Neurosci* 2004;20:658–70.
- [362] Bouvet JF, Delaleu JC, Holley A. The activity of olfactory receptor cells is affected by acetylcholine and substance P. *Neurosci Res* 1988;5:214–23.
- [363] Gutkowska J. Natriuretic peptides in the rat olfactory system. *Rev Bras Biol* 1996;36:79–88 Su 1 Pt 1.
- [364] Eisthen HL, Delay RJ, Wirsig-Wiechmann CR, Dionne VE. Neuromodulatory effects of gonadotropin releasing hormone on olfactory receptor neurons. *J Neurosci* 2000;20:3947–55.
- [365] Skipor J, Thiery JC. The choroid plexus-cerebrospinal fluid system: undervalued pathway of neuroendocrine signaling into the brain. *Acta Neurobiol Exp (Wars)* 2008;68:414–28.
- [366] Harrington MG, Fonteh AN, Oborina E, Liao P, Cowan RP, McComb G, Chavez JN, Rush J, Biringer RG, Huhmer AF. The morphology and biochemistry of nanostructures provide evidence for synthesis and signaling functions in human cerebrospinal fluid. *Cerebrospinal Fluid Res* 2009;6:10.
- [367] Swanson LW. Quest for the basic plan of nervous system circuitry. *Brain Res Rev* 2007;55:356–72.