Proofs as Programs Summer School
Eugene Oregon June - July 2002

Formalization in Coq

Formalizing an intuitionistic proof of the Fundamental Theorem of Algebra in Coq

Herman Geuvers, Freek Wiedijk
Jan Zwanenburg, Randy Pollack
Milad Niqui, Henk Barendregt

Fundamental Theorem of Algebra:

Every non-constant polynomial

$$f(z) = a_n z^n + a_{n-1} z^{n-1} + \ldots + a_1 z + a_0$$

with coefficients in $\mathbb{C}$ has a root
(a $z \in \mathbb{C}$ such that $f(z) = 0$).

- Overview / Motivation
- Constructive proof
- Axiomatic reasoning
- Completely formalized in the theorem prover Coq
- Some details of the proof
- Real numbers

Overview of the FTA project (Motivation)

- Formalize a large piece of real mathematics.
- Library for basic algebra and analysis, to be used by others
- Investigate the current limitations
- Try to manage this project. Three sequential/parallel phases:

Mathematical proof	LaTeX document (lots of details)
Theory development	Coq file (just defs and statements of lemmas)
Proof development	Coq file (proofs filled in)

  Try to keep these phases consistent!

- Constructive proof: reals are (potentially) infinite objects; algorithm.

Constructive proof

- Procedure for (always) finding a root.
- Equality on $\mathbb{R}$ is not decidable.
  (Not: $\forall x, y \in \mathbb{R} \quad (x = y) \vee (x \neq y)$)
- Constructively, a polynomial

$$f(z) = a_n z^n + a_{n-1} z^{n-1} + \ldots + a_1 z + a_0$$

  is non-constant if

  $a_k \# 0$ for some $k > 0$.

Axiomatic reasoning

- First define an algebraic hierarchy
  (semi-groups, monoids, groups, rings, fields, ordered fields)
- Advantages: Reuse of proven results; reuse of notation
- $\mathbb{R}$ is a Cauchy-complete Archimedean ordered field.
- $\mathbb{C} := \mathbb{R} \times \mathbb{R}$.
- A concrete instantiation for $\mathbb{R}$ has been constructed (Niqui)

Completely formalized in the theorem prover Coq

Sets and Basics	41 kb
Algebra (upto Ordered Fields)	165 kb
Reals	52 kb
Polynomials	113 kb
Real-valued functions / Basic Analysis	30 kb
Complex numbers	98 kb
FTA proof	70 kb
Construction of $\mathbb{R}$ (Niqui)	309 kb
Rational Tactic	49 kb

This is the size of our input files (definitions, lemmas, tactic scripts)

Some details of the proof

- Which proof to formalize?
- There are various constructive proofs of FTA:
  Weyl; Brouwer-de Loor; Bishop; Kneser (also in Troelstra and van Dalen).
- We have followed the Kneser proof.
- It can be seen as a constructive version of 'the' classical FTA proof.

The classical FTA proof

Suppose $|f(z)|$ is minimal with $|f(z)| \neq 0$.
We construct a z_0 with $|f(z_0)| < |f(z)|$.
We may assume that the minimum is reached for $z = 0$.

$$f(x) = a_0 + a_k x^k + O(x^{k+1})$$

with a_k the first coefficient that's not 0.
Now take

$$z_0 := \epsilon \sqrt[k]{-\frac{a_0}{a_k}}$$

with $\epsilon \in \mathbb{R}_{>0}$.
If ϵ is small enough, the part $O\left(z_0^{k+1}\right)$ will be negligible and we get a $z_0 \neq 0$ for which

$$|f(z_0)| = a_0 + a_k \left(\epsilon \sqrt[k]{-\frac{a_0}{a_k}}\right)^k = a_0(1 - \epsilon^k) < |f(0)|$$

The constructive FTA proof

Define an algorithm

Given $z \in \mathbb{C}$, construct a sequence $z, z_0, z_1, \ldots$ going to the root.

Problem: in the definition

$$z_0 := \epsilon \sqrt[k]{-\frac{a_0}{a_k}}$$

- ϵ must be small enough to neglect $O\left(z_0^{k+1}\right)$
- ϵ must be large enough to reach the root.

Solution: write

$$f(x) = a_0 + a_k x^k + \text{other terms}$$

and find k and z_0 such that $|a_k||z_0|^k$ is big enough w.r.t. the other terms and small enough compared to $|a_0|$.

Main Lemma

Given $|a_0| > 0$, $|a_1|, \ldots, |a_{n-1}| \geq 0$, $a_n = 1$, there are $r \in \mathbb{R}_{>0}$ and $k \in \{1, \ldots, n\}$ such that

$$\begin{aligned} r^n &< |a_0| \\ 3^{-2n^2}|a_0| &< |a_k|r^k < |a_0| \\ \Sigma_{i=1, i\neq k}^{n} |a_i| r^i &< (1 - 3^{-n})|a_k|r^k \end{aligned}$$

Now take

$$z := r \sqrt[k]{-\frac{a_0}{a_k} \cdot \frac{|a_k|}{|a_0|}}$$

So $a_k z^k$ points opposite to a_0. Then

$$|f(z)| < q \cdot |a_0|$$

with $q = 1 - 3^{-2n^2-n}$.

From this one constructs a sequence $(z_i)_{i\in\mathbb{N}}$ in $\mathbb{C}$ such that

$$\begin{aligned} |f(z_i)| &< q^i|a_0| \\ |z_{i+1} - z_i| &< (q^i|a_0|)^{1/n} \end{aligned}$$

This is a Cauchy sequence with limit a root of f.

Real Numbers in Coq:

- Axiomatic: a 'Real Number Structure' is a
  Cauchy-complete Archimedean ordered field.
- Prove FTA 'for all real numbers structures'.
- Construct a model to show that real number structures exist.
  (Cauchy sequences over an Arch. ordered field, say $\mathbb{Q}$)
- Prove that any two real number structures are isomorphic.

Axioms for Real Numbers:

- Algebraic hierarchy based on Constructive Setoids.
- Apartness $\#$ as basic

$$x = y \leftrightarrow \neg(x \# y)$$
$$x \# y \rightarrow (x \# z) \vee (y \# z)$$
$$x \# x$$
$$x \# y \rightarrow y \# x$$

- Field operation reciprocal

$$1/- : (\Sigma x{:}F.x \# 0) \rightarrow (\Sigma x{:}F.x \# 0)$$

- Cauchy sequences over Field F:
  g : nat $\rightarrow F$ is Cauchy if

$$\forall \epsilon{:}F_{>0}.\exists N{:}\mathbb{N}.\forall m \geq N(|g_m - g_N| < \epsilon)$$

The algebraic hierarchy of the FTA project

- In proving FTA, we have to deal with real numbers, complex numbers and polynomials
- Many of the properties we use are generic and algebraic.
- To be able to reuse results (also for future developments) we have defined a hierarchy of algebraic structures.
- Basic level: constructive setoids, $\langle A, \#_A, =_A \rangle$, with A : Set, $\#_A$ an apartness and $=_A$ an equivalence relation.
- Next level: semi-groups, $\langle S, + \rangle$, with S a setoid and $+$ an associative binary operation on S.

Inheritance via Coercions

We have the following coercions.

```
OrdField >-> Field >-> Ring >-> Group
                Group >-> Monoid >-> Semi_grp >-> Setoid
```

- All properties of groups are inherited by rings, fields, etc.
- Also notation is inherited:

  `x[+]y`

  denotes the addition of `x` and `y` for `x,y:G` from any semigroup (or monoid, group, ring,...) `G`.
- The coercions must form a tree, so there is no real multiple inheritance:
  E.g. it is not possible to define rings in such a way that it inherits both from its additive group and its multiplicative monoid.

Proofs by computation / Reflection

Needed for the proof of FTA:

Proofs of equalities between rational expressions like

$$\frac{1}{x+y} + \frac{1}{x-y} = \frac{2x}{x^2-y^2}$$

are obtained by partial reflection.

Following the reflection method:

$$[\![-]\!] : E \to \mathbb{R}$$

with E the type of rational expressions. So E contains a constructor `erecip : E -> E`

But in the case of rational expressions, the $[\![-]\!]$ can not be total.

$\rightsquigarrow$ partial reflection.

Axioms for Real Numbers ctd.:

- All Cauchy sequences have a limit:

$$\mathsf{SeqLim} : (\Sigma g{:}\mathsf{nat}{\to}F.\mathsf{Cauchy}\, g) \to F$$

$$\mathsf{CauchyProp} : \forall g{:}\mathsf{nat}{\to}F.(\mathsf{Cauchy}\, g) \to \forall \epsilon{:}F_{>0}.\exists N{:}\mathbb{N}.\forall m \geq N.(|g_m - (\mathsf{SeqLim}\, g)| < \epsilon)$$

- Axiom of Archimedes: (there are no non-standard elements)

$$\forall x{:}F.\exists n{:}\mathbb{N}(n > x)$$

NB: The axiom of Archimedes proves that 'ϵ-Cauchy sequences' and '$\frac{1}{k}$-Cauchy sequences' coincide (similar for limits):
Viz: $g : \mathsf{nat} \to F$ is a $\frac{1}{k}$-Cauchy sequence if

$$\forall k{:}\mathbb{N}.\exists N{:}\mathbb{N}.\forall m \geq N(|g_m - g_N| < \frac{1}{k+1})$$

Construction of a Real Number Structure:

- Construct $\mathbb{Q} := \{\langle p, n\rangle \mid p{:}\mathbb{Z}, n{:}\mathbb{N}\}$ with intended interpretation $\langle p, n\rangle \mapsto \frac{p}{n+1}$.
- Define all operations and relations, turning $\mathbb{Q}$ into an Archimedean constructive ordered field.
- For F an (Archimedean) constructive ordered field, define the Cauchy completion of F, CSeq_F.

$$h < g := \exists \epsilon : F_{>0}.\exists N{:}\mathbb{N}\forall m \geq N.(g_m - h_m > \epsilon).$$

$$g_m^{-1} := \begin{cases} 0 & \text{if } \; m < N \\ \frac{1}{g_m} & \text{if } \; m \geq N \end{cases}$$

where N is such that $\forall m \geq N.|g_m| > \epsilon$ for some $\epsilon > 0$.

- Prove that CSeq_F is a Real Number Structure.

Categoricity of Real Number Structures:

- A morphism between R_1 and R_2 is a $\varphi : R_1 \to R_2$ that
  1. is strongly extensional: $\forall x, y : R_1.\ \varphi(x) \# \varphi(y) \to x \# y$.
  2. preserves order, addition and multiplication.
- A morphism preserves 'all' structure (Cauchy property, limits, ...)
  (Use: $\forall y \in R_2 . \exists x \in R_1 (\varphi(x) < y)$.)
- $R_1 \simeq R_2$ if $\varphi : R_1 \to R_2$ for some bijective morphism φ.

Categoricity of Real Number Structures ctd.:

- All Real Number Structures are isomorphic to $\mathbb{Q}^{\mathbb{N}}$:



- $\mathsf{Inj}_R : \mathbb{Q} \to R$ canonically maps rationals into R.
- $\mathsf{Seq} : R \to \mathbb{Q}^{\mathbb{N}}$: For $x : R$, define the $\mathbb{Q}$-sequences $\{q_i\}_{i=0}^{\infty}$ (increasing) and $\{r_i\}_{i=0}^{\infty}$ (decreasing), such that
  1. $\forall n{:}\mathbb{N}(r_n - q_n = (r_0 - q_0)(\frac{2}{3})^n)$
  2. $\forall n{:}\mathbb{N}(\mathsf{Inj}_R(q_n) \leq x \leq \mathsf{Inj}_R(r_n))$

Some Conclusions:

- Real mathematics, involving both a bit of algebra and a bit of analysis can be formalised completely within a theorem prover (Coq).
- Setting up a basic library and some good proof automation procedures is a substantial part of the work.
- Computationally, the behaviour of the algorithm depends mainly on the representation of the reals.
- The non-determinism in the root-finding algorithm (non-continuity of the root-finding function) lies in the taking of a k-th root in $\mathbb{C}$.
- How to present a proof?? Curry Howard: proofs-as-terms Converse??

Some Conclusions:

- Real mathematics, involving both a bit of algebra and a bit of analysis can be formalised completely within a theorem prover (Coq).
- Setting up a basic library and some good proof automation procedures is a substantial part of the work.
- Computationally, the behaviour of the algorithm depends mainly on the representation of the reals.
- The non-determinism in the root-finding algorithm (non-continuity of the root-finding function) lies in the taking of a k-th root in $\mathbb{C}$.
- How to present a proof?? Curry-Howard: proofs-as-terms
  Converse: Hurry-Coward: present terms as 'readable' proofs