


# Markov Independence




# A Bayesian network



Thus:  $P(\text{FL}, \text{MY}, \text{FE}) = P(\text{MY} | \text{FL}, \text{FE}) P(\text{FE} | \text{FL}) P(\text{FL})$

Example:  $P(\neg fl, my, fe) = 0.20 \cdot 0.1 \cdot 0.9 = 0.018$

# Independence and reasoning




# Independence and reasoning

Conclusion: the arc from FEVER to MYALGIA can be removed, and hence only

$$P(\text{MY} | \text{FL}) (= P(\text{MY} | \text{FL}, \text{FE}))$$

need be specified




# Independence relation

Let  $X, Y, Z \subseteq V$  be sets of (random) variables, and let  $P$  be a probability distribution of  $V$  then  $X$  is called **conditionally independent** of  $Y$  given  $Z$ , denoted as

$$X \perp\!\!\!\perp_P Y \mid Z, \text{ iff } P(X \mid Y, Z) = P(X \mid Z)$$

**Note:** This relation is completely defined in terms of the probability distribution  $P$ , but there is a *relationship to graphs*, for example:



# Equivalences

The following conditions are equivalent:

- $P(X \mid Y, Z) = P(X \mid Z)$  if  $P(Y, Z) > 0$
- $P(X, Y \mid Z) = P(X \mid Z)P(Y \mid Z)$  if  $P(Y, Z) > 0$
- $P(X \mid Y, Z)$  can be represented as the real function  $\psi(X, Z)$ , called a **potential**
- $P(X, Y \mid Z)$  can be written as  $\phi(X, Z)\psi(Y, Z)$ , with real potential functions  $\phi$  and  $\psi$
- $P(X, Y, Z) = P(X \mid Z)P(Y \mid Z)P(Z)$
- $P(X, Y, Z) = P(X, Z)P(Y, Z)/P(Z)$  if  $P(Z) > 0$

N.B. potentials are non-negative real functions, very similar to probability distributions, but they need not be normalised

## Properties of the $\perp\!\!\!\perp_P$ relation (1)

**P1 Symmetry:** Let  $X, Y, Z \subseteq V$  be sets of variables, then:

$$X \perp\!\!\!\perp_P Y \mid Z \iff Y \perp\!\!\!\perp_P X \mid Z$$

**Proof:**  $X \perp\!\!\!\perp_P Y \mid Z \iff P(X \mid Y, Z) = P(X \mid Z)$  (1)

$$\begin{aligned} \frac{P(X, Y \mid Z)}{P(Y \mid Z)} &= \frac{P(Y \mid X, Z)P(X \mid Z)}{P(Y \mid Z)} \\ \Rightarrow P(Y \mid X, Z) &= \frac{P(X, Y \mid Z)}{P(X \mid Z)} \\ &= \frac{P(X \mid Y, Z)P(Y \mid Z)}{P(X \mid Z)} \\ &\stackrel{(1)}{=} P(Y \mid Z) \\ &\iff Y \perp\!\!\!\perp_P X \mid Z \end{aligned}$$

## Properties of the $\perp\!\!\!\perp_P$ relation (2)

**P2 Decomposition:** Let  $X, Y, Z \subseteq V$  be sets of variables with  $U \subseteq X$  then:

$$X \perp\!\!\!\perp_P Y \mid Z \Rightarrow U \perp\!\!\!\perp_P Y \mid Z$$

**Proof:**  $X \perp\!\!\!\perp_P Y \mid Z \iff P(X \mid Y, Z) = P(X \mid Z)$  (1)

$$\begin{aligned} P(U \mid Y, Z) &= \sum_{X \setminus U} P(U, X \mid Y, Z) \\ &= \sum_{X \setminus U} P(X \mid Y, Z), \text{ as } U \subseteq X \\ &\stackrel{(1)}{=} \sum_{X \setminus U} P(X \mid Z) \\ &= P(U \mid Z) \iff U \perp\!\!\!\perp_P Y \mid Z \end{aligned}$$

## Properties of the $\perp\!\!\!\perp_P$ relation (3)

**P3 Weak union:** Let  $X, Y, Z, U \subseteq V$  be sets of variables with  $U \subseteq X$ , then

$$X \perp\!\!\!\perp_P Y \mid Z \Rightarrow X \perp\!\!\!\perp_P Y \mid Z \cup U$$

**Proof:** ... *DIY*

**P4 Contraction:** Let  $X, Y, Z, W \subseteq V$  be sets of variables, then:

$$\begin{aligned} X \perp\!\!\!\perp_P Y \mid Z \wedge X \perp\!\!\!\perp_P W \mid Y \cup Z \\ \Rightarrow X \perp\!\!\!\perp_P W \cup Y \mid Z \end{aligned}$$

**Proof:** ... *DIY*

Any model that offers a representation of axioms **(P1)** to **(P4)** is called a **semi-graphoid**

Lecture 4: Independence – p. 9/4

## Properties of the $\perp\!\!\!\perp_P$ relation (4)

**P5 Intersection:** Let  $X, Y, Z, W \subseteq V$  be sets of variables, then


$$\begin{aligned} X \perp\!\!\!\perp_P Y \mid Z \cup W \wedge X \perp\!\!\!\perp_P Z \mid Y \cup W \\ \Rightarrow X \perp\!\!\!\perp_P Y \cup Z \mid W \end{aligned}$$

**Proof:** ... *DIY*

- This axiom only holds for *strictly positive* probability distributions, i.e. probability distributions that do not represent *logical relationships*.
- Any model that satisfies axioms **(P1)** to **(P5)** is called a **graphoid**


Lecture 4: Independence – p. 10/4

## From probabilities to independence relation



Lecture 4: Independence – p. 11/4

## Definition of an independence relation



Lecture 4: Independence – p. 12/4

## Definition of an independence relation

Let  $X, Y, Z, W \subseteq V$  be sets of objects. The **independence relation**  $\perp\!\!\!\perp \subseteq \wp(V) \times \wp(V) \times \wp(V)$  is defined such that the following properties hold:

- **Symmetry:**  $X \perp\!\!\!\perp Y \mid Z \iff Y \perp\!\!\!\perp X \mid Z$
- **Decomposition:**  $X \perp\!\!\!\perp Y \mid Z \Rightarrow U \perp\!\!\!\perp Y \mid Z$ , with  $U \subseteq X$
- **Weak union:**  $X \perp\!\!\!\perp Y \mid Z \Rightarrow X \perp\!\!\!\perp Y \mid Z \cup U$ , with  $U \subseteq X$
- **Contraction:**  
 $X \perp\!\!\!\perp Y \mid Z \wedge X \perp\!\!\!\perp W \mid Y \cup Z \Rightarrow X \perp\!\!\!\perp W \cup Y \mid Z$

i.e.  $\perp\!\!\!\perp$  defines a **semi-graphoid**. Note that the intersection property need not hold

Lecture 4: Independence – p. 13/4

## How to define an independence relation?

- List all the instances of  $\perp\!\!\!\perp$
- List some of the instances of  $\perp\!\!\!\perp$  and add axioms from which other instances can be derived
- Define a joint probability distribution  $P$  and look into the numbers to see which instances of the independence relation  $\perp\!\!\!\perp$  hold (this yields  $\perp\!\!\!\perp_P$ )
- **Use a graph** to encode  $\perp\!\!\!\perp$ , which yields  $\perp\!\!\!\perp_G$  (so, what type of graph — directed, undirected, chain?)

Lecture 4: Independence – p. 15/4

## Use of independence axioms

**Lemma** Let  $X, Y, Z, W \subseteq V$  be sets of objects, then

$$X \perp\!\!\!\perp Y \mid Z \wedge X \cup Z \perp\!\!\!\perp W \mid Y \\ \Rightarrow X \perp\!\!\!\perp W \mid Z$$

**Proof:** It holds that

$$X \cup Z \perp\!\!\!\perp W \mid Y \Rightarrow_{\text{symm}} W \perp\!\!\!\perp X \cup Z \mid Y \\ \Rightarrow_{\text{wu}} W \perp\!\!\!\perp X \mid Y \cup Z \Rightarrow_{\text{symm}} X \perp\!\!\!\perp W \mid Y \cup Z$$

From  $X \perp\!\!\!\perp Y \mid Z$  and  $X \perp\!\!\!\perp W \mid Y \cup Z$ , using contraction, it follows that  $X \perp\!\!\!\perp W \cup Y \mid Z$ . Now, by using decomposition, it follows that  $X \perp\!\!\!\perp W \mid Z$

Lecture 4: Independence – p. 14/4


## Explicit enumeration

Consider  $V = \{1, 2, 3, 4\}$  and  $\perp\!\!\!\perp$ :

$\{1\} \perp\!\!\!\perp \{4\} \mid \emptyset$	$\{4\} \perp\!\!\!\perp \{2\} \mid \{1\}$	$\{2\} \perp\!\!\!\perp \{4\} \mid \emptyset$
$\{4\} \perp\!\!\!\perp \{3\} \mid \{1\}$	$\{3\} \perp\!\!\!\perp \{4\} \mid \emptyset$	$\{4\} \perp\!\!\!\perp \{2, 3\} \mid \{1\}$
$\{4\} \perp\!\!\!\perp \{1\} \mid \emptyset$	$\{1\} \perp\!\!\!\perp \{4\} \mid \{2\}$	$\{4\} \perp\!\!\!\perp \{2\} \mid \emptyset$
$\{3\} \perp\!\!\!\perp \{4\} \mid \{2\}$	$\{4\} \perp\!\!\!\perp \{3\} \mid \emptyset$	$\{1, 3\} \perp\!\!\!\perp \{4\} \mid \{2\}$
$\{1, 2\} \perp\!\!\!\perp \{4\} \mid \emptyset$	$\{4\} \perp\!\!\!\perp \{1\} \mid \{2\}$	$\{1, 3\} \perp\!\!\!\perp \{4\} \mid \emptyset$
$\{4\} \perp\!\!\!\perp \{3\} \mid \{2\}$	$\{2, 3\} \perp\!\!\!\perp \{4\} \mid \emptyset$	$\{4\} \perp\!\!\!\perp \{1, 3\} \mid \{2\}$
$\{4\} \perp\!\!\!\perp \{1, 2\} \mid \emptyset$	$\{1\} \perp\!\!\!\perp \{4\} \mid \{3\}$	$\{4\} \perp\!\!\!\perp \{1, 3\} \mid \emptyset$
$\{2\} \perp\!\!\!\perp \{4\} \mid \{3\}$	$\{4\} \perp\!\!\!\perp \{2, 3\} \mid \emptyset$	$\{1, 2\} \perp\!\!\!\perp \{4\} \mid \{3\}$
$\{1, 2, 3\} \perp\!\!\!\perp \{4\} \mid \emptyset$	$\{1\} \perp\!\!\!\perp \{2\} \mid \{4\}$	$\{4\} \perp\!\!\!\perp \{1, 2, 3\} \mid \emptyset$
$\{2\} \perp\!\!\!\perp \{1\} \mid \{4\}$	$\{1\} \perp\!\!\!\perp \{2\} \mid \emptyset$	$\{3\} \perp\!\!\!\perp \{4\} \mid \{1, 2\}$
$\{2\} \perp\!\!\!\perp \{1\} \mid \emptyset$	$\{4\} \perp\!\!\!\perp \{3\} \mid \{1, 2\}$	$\{1, 4\} \perp\!\!\!\perp \{2\} \mid \emptyset$
$\{2\} \perp\!\!\!\perp \{4\} \mid \{1, 3\}$	$\{2, 4\} \perp\!\!\!\perp \{1\} \mid \emptyset$	$\{4\} \perp\!\!\!\perp \{2\} \mid \{1, 3\}$
$\{2\} \perp\!\!\!\perp \{1, 4\} \mid \emptyset$	$\{1\} \perp\!\!\!\perp \{4\} \mid \{2, 3\}$	$\{1\} \perp\!\!\!\perp \{2, 4\} \mid \emptyset$
$\{4\} \perp\!\!\!\perp \{1\} \mid \{2, 3\}$	$\{2\} \perp\!\!\!\perp \{4\} \mid \{1\}$	$\{4\} \perp\!\!\!\perp \{1, 2\} \mid \{3\}$
$\{3\} \perp\!\!\!\perp \{4\} \mid \{1\}$	$\{4\} \perp\!\!\!\perp \{1\} \mid \{3\}$	$\{2, 3\} \perp\!\!\!\perp \{4\} \mid \{1\}$
$\{4\} \perp\!\!\!\perp \{2\} \mid \{3\}$		

Lecture 4: Independence – p. 16/4

## As an undirected graph




Basic idea:

- Each variable  $V$  is represented as a vertex in an undirected graph  $G = (V(G), E(G))$ , with set of vertices  $V(G)$  and set of edges  $E(G)$
- the **independence relation**  $\perp\!\!\!\perp_G$  is encoded as the **absence of edges**; a missing edge between vertices  $u$  and  $v$  indicates that random variables  $X_u$  and  $X_v$  are (conditionally) independent

Lecture 4: Independence – p. 17/4

## Example

Consider the following undirected graph  $G$ :



- $\{1\} \perp\!\!\!\perp_G \{3, 6\} \mid \{2\}$
- $\{4\} \perp\!\!\!\perp_G \{6\} \mid \{2, 5\}$
- $\{4\} \perp\!\!\!\perp_G \{6\} \mid \{1, 2, 3, 5\}$
- $\{1\} \not\perp\!\!\!\perp_G \{5\} \mid \{4\}$ , as the path  $1 - 2 - 5$  does not contain 4
- $\{1, 5, 6\} \perp\!\!\!\perp_G \{7\} \mid \emptyset$

Lecture 4: Independence – p. 19/4

## Global Markov property – separation

Let  $G = (V(G), E(G))$  be an undirected graph, and let  $U, Z, W \subseteq V(G)$  be sets of vertices in  $G$ . The set  $W$  **(u-)separates**  $U$  and  $Z$ , denoted as

$$U \perp\!\!\!\perp_G Z \mid W$$

if every path from a vertex in  $U$  to a vertex in  $Z$  contains at least one vertex in  $W$ ; otherwise these sets are **(u-)connected**

Remarks:

- This criterion is known as the **global Markov property** or **(u-)separation criterion** for undirected graphs
- Note that  $\perp\!\!\!\perp_G$  indicates that the independence relation is defined in terms of  $G$  (cf.  $\perp\!\!\!\perp_P$ )
- If there are no paths between two vertices  $u$  and  $v$ , then  $\{u\} \perp\!\!\!\perp_G \{v\} \mid \emptyset$

Lecture 4: Independence – p. 18/4

## D-map and I-map

Let  $V$  be a set and let  $\perp\!\!\!\perp$  be an independence relation defined on  $V$ . Let  $G = (V(G), E(G))$  be an undirected graph with  $V(G) = V$ , then for each  $X, Y, Z \subseteq V$ :

- $G$  is called an undirected **dependence map**, **D-map** for short, if

$$X \perp\!\!\!\perp Y \mid Z \Rightarrow X \perp\!\!\!\perp_G Y \mid Z$$

- $G$  is called an undirected **independence map**, **I-map** for short, if

$$X \perp\!\!\!\perp_G Y \mid Z \Rightarrow X \perp\!\!\!\perp Y \mid Z$$

- $G$  is called an undirected **perfect map**, or **P-map** for short, if  $G$  is both a D-map and an I-map, or, equivalently

$$X \perp\!\!\!\perp Y \mid Z \iff X \perp\!\!\!\perp_G Y \mid Z$$

Lecture 4: Independence – p. 20/4

## D-map and I-map for $\perp\!\!\!\perp_P$

Let  $P$  be probability distribution of  $X$ . Let  $G = (V(G), E(G))$  be an undirected graph, then for each  $U, W, Z \subseteq V(G)$ :

- $G$  is called an undirected **dependence map**, **D-map** for short, if

$$X_U \perp\!\!\!\perp_P X_W \mid X_Z \Rightarrow U \perp\!\!\!\perp_G W \mid Z$$

- $G$  is called an undirected **independence map**, **I-map** for short, if

$$U \perp\!\!\!\perp_G W \mid Z \Rightarrow X_U \perp\!\!\!\perp X_W \mid X_Z$$

- $G$  is called an undirected **perfect map**, or **P-map** for short, if  $G$  is both a D-map and an I-map, or, equivalently

$$X_U \perp\!\!\!\perp_P X_W \mid X_Z \iff U \perp\!\!\!\perp_G W \mid Z$$

Lecture 4: Independence – p. 21/4


## Markov network

A pair  $\mathcal{M} = (G, P)$ , where

- $G = (V(G), E(G))$  is an *undirected* graph with set of vertices  $V(G)$  and set of edges  $E(G)$ ,
- $P$  is a joint probability distribution of  $X_{V(G)}$ , and
- $G$  is an *I-map* of  $P$

is said to be a **Markov network** or **Markov random field**

**Example**  $\mathcal{M} = (G, \phi) = (G, P)$ :



Potential:

$$\phi(X_1, X_2, X_3) = \psi(X_1, X_2)\tau(X_2, X_3),$$

or joint probability distribution:

$$P(X_1, X_2, X_3) = \frac{P(X_1, X_2)P(X_2, X_3)}{P(X_2)}$$

Lecture 4: Independence – p. 23/4


## Examples D-maps

Let  $V = \{1, 2, 3, 4\}$  be a set and  $X_V$  the corresponding set of random variables, and consider the independence relation  $\perp\!\!\!\perp_P$ , defined by

$$\{X_1\} \perp\!\!\!\perp_P \{X_4\} \mid \{X_2, X_3\}$$

$$\{X_2\} \perp\!\!\!\perp_P \{X_3\} \mid \{X_1, X_4\}$$

The following undirected graphs are examples of D-maps:



Lecture 4: Independence – p. 22/4

## D-maps and I-maps again

Let  $\perp\!\!\!\perp$  be an independence relation. D-maps and I-maps are limited in expressiveness in the following sense:

- A pair of neighbour vertices in a D-map for  $\perp\!\!\!\perp$  are dependent. However, not all dependent variables are neighbours
- A pair of non-neighbour variables in an I-map for  $\perp\!\!\!\perp$  corresponds to independent variables, but not each pair of independent variables in an I-map are non-neighbours


Lecture 4: Independence – p. 24/4

## Examples of I-maps

Let  $V = \{1, 2, 3, 4\}$  be a set with random variables  $X_V$ , and consider the independence relation  $\perp\!\!\!\perp_P$ :

$$\begin{aligned} \{X_1\} &\perp\!\!\!\perp_P \{X_4\} \mid \{X_2, X_3\} \\ \{X_2\} &\perp\!\!\!\perp_P \{X_3\} \mid \{X_1, X_4\} \end{aligned}$$

The following undirected graphs are examples of I-maps:



(So, what is the P-map?)

Lecture 4: Independence – p. 25/4

## Obvious properties

**Lemma** For each independence relation  $\perp\!\!\!\perp$  there exists an undirected D-map.

**Proof:**

The undirected graph  $G = (V, \emptyset)$  is a D-map for  $\perp\!\!\!\perp$ .

**Lemma** For each independence relation  $\perp\!\!\!\perp$  there exists an undirected I-map.


**Proof:**

The undirected graph  $G = (V, V \times V)$  is an I-map for  $\perp\!\!\!\perp$ .

Lecture 4: Independence – p. 26/4

## Expressiveness: directed vs undirected

**Directed graphs** are more subtle when it comes to expressing independence information than **undirected graphs**



vs

Lecture 4: Independence – p. 27/4

## d-Separation: 3 situations

A **chain  $k$**  (= path in undirected underlying graph) in an acyclic directed graph  $G = (V(G), A(G))$  can be **blocked**:

**Diverging**




2 **blocks** (d-separates) 1 and 3:  $\{1\} \perp\!\!\!\perp \{3\} \mid \{2\}$

**Serial**



2 **blocks** (d-separates) 1 and 3:  $\{1\} \perp\!\!\!\perp \{3\} \mid \{2\}$

**Converging**




2 **d-connects** 1 and 3:  $\{1\} \not\perp\!\!\!\perp \{3\} \mid \{2\}$

(same holds for successors of 2); note  $\{1\} \perp\!\!\!\perp \{3\} \mid \emptyset$

Lecture 4: Independence – p. 28/4

## Example blockage



- The chain 4, 2, 5 from 4 to 5 is blocked by {2}
- The chain 1, 2, 5, 6 from 1 to 6 is blocked by {5}, and also by {2} and {2, 5}
- The chain 3, 4, 6, 5 from 3 to 5 is blocked by {4} and {4, 6}, but *not* by {6}

Lecture 4: Independence – p. 29/4

## Directed global Markov property

Let  $G = (V(G), A(G))$  be an acyclic directed graph, and let  $U, W, Z \subseteq V(G)$  be sets of vertices in  $G$ . The set  $Z$  **d-separates**  $U$  and  $W$ , denoted as

$$U \perp\!\!\!\perp_G^d W \mid Z$$

if every chain from a vertex in  $U$  to a vertex in  $W$  is blocked by  $Z$

### Remarks

- This criterion is known as the **global Markov property** or **d-separation criterion** for acyclic directed graphs
- Note that  $\perp\!\!\!\perp_G^d$  indicates that the independence relation is defined in terms of  $G$  (cf.  $\perp\!\!\!\perp_P$ )


Lecture 4: Independence – p. 30/4

## There is also a local Markov property

Let  $G = (V(G), A(G))$  be an acyclic, directed graph, then the following **local Markov property** holds:

$$\{v_i\} \perp\!\!\!\perp_G^d \nu(v_i) \mid \pi(v_i)$$

with  $\nu(v_i)$  *non-descendants* of vertex  $v_i$ , and  $\pi(v_i)$  set of parents



Lecture 4: Independence – p. 31/4

## Directed D-map and I-map

Let  $V$  be a set and let  $\perp\!\!\!\perp$  be an independence relation defined on  $V$ . Let  $G = (V(G), A(G))$  be an acyclic directed graph, then for each  $X, Y, Z \subseteq V$ :

- $G$  is called a directed **dependence map**, **D-map** for short, if

$$X \perp\!\!\!\perp Y \mid Z \Rightarrow X \perp\!\!\!\perp_G^d Y \mid Z$$

- $G$  is called a directed **independence map**, **I-map** for short, if

$$X \perp\!\!\!\perp_G^d Y \mid Z \Rightarrow X \perp\!\!\!\perp Y \mid Z$$

- $G$  is called a directed **perfect map**, or **P-map** for short, if  $G$  is both a D-map and an I-map, or, equivalently

$$X \perp\!\!\!\perp Y \mid Z \iff X \perp\!\!\!\perp_G^d Y \mid Z$$


Lecture 4: Independence – p. 32/4

## Examples directed I-maps

Consider the following independence relation  $\perp\!\!\!\perp_P$ :

$$\begin{aligned} \{X_1\} &\perp\!\!\!\perp_P \{X_2\} \mid \emptyset \\ \{X_1, X_2\} &\perp\!\!\!\perp_P \{X_4\} \mid \{X_3\} \end{aligned}$$

and the following directed I-maps of  $P$ :



Lecture 4: Independence – p. 33/4


## Minimal directed I-map

In the context of Bayesian networks, we are interested in I-maps that contain as few arcs as possible (makes probability tables smaller), i.e. **minimal directed I-maps**

Let  $G = (V(G), A(G))$  be an acyclic directed graph and let  $P(X_{V(G)})$  be a probability distribution of  $X_{V(G)}$ .  $G$  is said to be a **minimal directed I-map** of  $P$ , if

- $G$  is a directed I-map of  $P$ , and
- none of the subgraphs of  $G$  is a directed I-map of  $P$

Example:



Lecture 4: Independence – p. 34/4

## Relationship directed and undirected graphs

- Directed graphs contain independences that become dependences after conditioning (instantiating variables)
- Undirected graphs do not have this property
- However, undirected subgraphs can be generated, by making potentially dependent parents of a child dependent

Example:




Lecture 4: Independence – p. 35/4

## Moralisation

Let  $G$  be an acyclic directed graph; its associated undirected **moral graph**  $G^m$  can be constructed by **moralisation**:

1. add lines to all non-connected vertices, which have a common child, or descendant of a common child, and
2. replace each arc with a line in the resulting graph




Lecture 4: Independence – p. 36/4

# Moralisation

Let  $G$  be an acyclic directed graph; its associated undirected **moral graph**  $G^m$  can be constructed by **moralisation**:

1. add lines to all non-connected vertices, which have a common child, or descendant of a common child, and
2. replace each arc with a line in the resulting graph



Lecture 4: Independence – p. 36/4

# Comments

- Resulting undirected (moral) graph is an I-map of the associated probability distribution
- However, it contains **too many dependences!**

Example:  $\{1\} \perp\!\!\!\perp_G^d \{3\} \mid \emptyset$ , whereas  $\{1\} \not\perp\!\!\!\perp_{G^m} \{3\} \mid \emptyset$




- Conclusion: make moralisation **'dynamic'** (i.e. a function of the set on which we condition)
- For this the notion of 'ancestral set' is required

Lecture 4: Independence – p. 37/4

# Ancestral set

Let  $G = (V(G), A(G))$  be an acyclic directed graph, then if for  $W \subseteq V(G)$  it holds that  $\pi(v) \subseteq W$  for all  $v \in W$ , then  $W$  is called an **ancestral set** of  $W$ .  $An(W)$  denotes the **smallest** ancestral set containing  $W$



Lecture 4: Independence – p. 38/4

# 'Dynamic' moralisation


Let  $P$  be a joint probability distribution of a Bayesian network  $\mathcal{B} = (G, P)$ , then

$$X_U \perp\!\!\!\perp_P X_V \mid X_W$$

holds iff  $U$  and  $V$  are (u-)separated by  $W$  in the moral induced subgraph  $G^m$  of  $G$  with vertices  $An(U \cup V \cup W)$

Example:

$$X_1 \not\perp\!\!\!\perp_P X_3 \mid X_2; \quad An(\{1, 2, 3\}) = \{1, 2, 3\}$$



Lecture 4: Independence – p. 39/4

## 'Dynamic' moralisation


Let  $P$  be a joint probability distribution of a Bayesian network  $\mathcal{B} = (G, P)$ , then

$$X_U \perp\!\!\!\perp_P X_V \mid X_W$$

holds iff  $U$  and  $V$  are (u-)separated by  $W$  in the moral induced subgraph  $G^m$  of  $G$  with vertices  $An(U \cup V \cup W)$

Example:

$$X_1 \perp\!\!\!\perp_P X_3 \mid \emptyset; \quad An(\{1, 3\}) = \{1, 3\}$$



Lecture 4: Independence – p. 39/4

## Moralisation and d-separation

Let  $G = (V(G), A(G))$  be an acyclic directed graph and let  $U, W, S \subseteq V(G)$  be disjoint sets of vertices. Then,  $U$  and  $W$  are d-separated by  $S$ , i.e.

$$U \perp\!\!\!\perp_G^d W \mid S$$

iff  $U$  and  $W$  are separated in the moral graph of the set of vertices  $An(U \cup W \cup S)$ , i.e.


$$U \perp\!\!\!\perp_{G_{An(U \cup W \cup S)}^m} W \mid S$$

**Proof:** Cowell et al, "Probabilistic Networks and Expert Systems", 1999, Springer, New York, page 72

Lecture 4: Independence – p. 40/4

## Example (1)


$$\{10\} \not\perp\!\!\!\perp_G^d \{13\} \mid \{7, 8\}$$



Lecture 4: Independence – p. 41/4

## Example (1)


$$\{10\} \not\perp\!\!\!\perp_{G_{An(\{10,7,8,13\})}^m} \{13\} \mid \{7, 8\}$$



Lecture 4: Independence – p. 41/4

## Example (2)

$$\{10\} \perp\!\!\!\perp_G^d \{13\} \mid \emptyset$$



Lecture 4: Independence – p. 42/4

## Example (2)

$$\{10\} \not\perp\!\!\!\perp_{G_{\text{An}(\{10,13\})}^m} \{13\} \mid \emptyset$$



Lecture 4: Independence – p. 42/4

## Conclusions

- Conditional independence is defined as a logic that supports:
  - symbolic reasoning about dependence and independence information
  - makes it possible to abstract away from the numerical detail of probability distributions
  - the process of assessing probability distributions
- Looking at graphs makes it easier to find probability distributions that are **equivalent** (important in learning)
- **Conditional** independence is currently being extended towards **causal** independence (a logic of causality) = **maximal ancestral graphs**

Lecture 4: Independence – p. 43/4