

Matrix Calculations: Vector Spaces

A. Kissinger

Institute for Computing and Information Sciences
Radboud University Nijmegen

Version: autumn 2017



Outline

Non-homogeneous systems

Vector spaces



From last time

Homogeneous systems have 0's in the RHS of all equations.

Given a homogeneous system in n variables:

- A **basic solution** is a **non-zero** solution of the system.
- If there are n pivots in its Echelon form, $\mathbf{0} = (0, \dots, 0)$ is the **unique solution**, so no basic solutions.
- If there are $p < n$ pivots in its Echelon form, it has $n - p$ **linearly independent** basic solutions.
- Two methods for finding them: plugging in free variables or deleting non-pivot columns, one-by-one



Non-homogeneous case: subtracting solutions

Theorem

For two solutions \mathbf{s} and \mathbf{p} of a **non-homogeneous** system of equations, the difference $\mathbf{s} - \mathbf{p}$ is a solution of the associated **homogeneous** system.

Proof: Let $a_1x_1 + \dots + a_nx_n = b$ be an equation in the non-homogeneous system. Then:

$$\begin{aligned} & a_1(s_1 - p_1) + \dots + a_n(s_n - p_n) \\ &= \left(a_1s_1 - a_1p_1 \right) + \dots + \left(a_ns_n - a_np_n \right) \\ &= \left(a_1s_1 + \dots + a_ns_n \right) - \left(a_1p_1 + \dots + a_np_n \right) \\ &= b - b \quad \text{since the } \mathbf{s} \text{ and } \mathbf{p} \text{ are solutions} \\ &= 0. \end{aligned}$$

General solution for non-homogeneous systems


Theorem

Assume a non-homogeneous system has a solution given by the vector \mathbf{p} , which we call a *particular solution*.

Then any other solution \mathbf{s} of the non-homogeneous system can be written as

$$\mathbf{s} = \mathbf{p} + \mathbf{h}$$

where \mathbf{h} is a solution of the associated homogeneous system.

Proof: Let \mathbf{s} be a solution of the non-homogeneous system. Then $\mathbf{h} = \mathbf{s} - \mathbf{p}$ is a solution of the associated homogeneous system. Hence we can write \mathbf{s} as $\mathbf{p} + \mathbf{h}$, for \mathbf{h} some solution of the associated homogeneous system. 

Example: solutions of a non-homogeneous system

- Consider the non-homogeneous system
$$\begin{cases} x + y + 2z = 9 \\ y - 3z = 4 \end{cases}$$
- with solutions: $(0, 7, 1)$ and $(5, 4, 0)$
- We can write $(0, 7, 1)$ as: $(5, 4, 0) + (-5, 3, 1)$
- where:
 - $\mathbf{p} = (5, 4, 0)$ is a **particular solution** (of the original system)
 - $(-5, 3, 1)$ is a solution of the associated **homogeneous** system:
$$\begin{cases} x + y + 2z = 0 \\ y - 3z = 0 \end{cases}$$
- Similarly, $(10, 1, -1)$ is a solution of the non-homogeneous system and
$$(10, 1, -1) = (5, 4, 0) + (5, -3, -1)$$
- where:
 - $(5, -3, -1)$ is a solution of the associated **homogeneous** system.

General solution for non-homogeneous systems, concretely

Theorem

The general solution of a non-homogeneous system of equations in n variables is given by a *parametrization* as follows:

$$(s_1, \dots, s_n) = (p_1, \dots, p_n) + c_1(v_{11}, \dots, v_{1n}) + \dots + c_k(v_{k1}, \dots, v_{kn})$$

for $c_1, \dots, c_k \in \mathbb{R}$,

where

- (p_1, \dots, p_n) is a *particular solution*
- $(v_{11}, \dots, v_{1n}), \dots, (v_{k1}, \dots, v_{kn})$ are *basic solutions* of the associated homogeneous system.
- So $c_1(v_{11}, \dots, v_{1n}) + \dots + c_k(v_{k1}, \dots, v_{kn})$ is a general solution for the associated homogeneous system.

Finding a particular solution

- **Recall:** we found basic solutions by setting **all but one** of the free variables to zero and solving the **homogeneous system**
- To find a particular solution, set **all** the free variables to zero and solving the **non-homogeneous system**
- In other words, remove **all** the non-pivot columns:

$$\left(\begin{array}{ccccc|c} \boxed{1} & 1 & 1 & 1 & 1 & 3 \\ 0 & 0 & \boxed{1} & 2 & 3 & 1 \\ 0 & 0 & 0 & 0 & \boxed{1} & 4 \end{array} \right) \mapsto \left(\begin{array}{ccc|c} \boxed{1} & 1 & 1 & 3 \\ 0 & \boxed{1} & 3 & 1 \\ 0 & 0 & \boxed{1} & 4 \end{array} \right)$$

- Solve. Then, add zeros back in for the free variables:

$$(10, -11, 4) \mapsto (10, 0, -11, 0, 4)$$

Elaborated example, part I

- Consider the **non-homogeneous** system of equations given by the augmented matrix in echelon form:

$$\left(\begin{array}{ccccc|c} \boxed{1} & 1 & 1 & 1 & 1 & 3 \\ 0 & 0 & \boxed{1} & 2 & 3 & 1 \\ 0 & 0 & 0 & 0 & \boxed{1} & 4 \end{array} \right)$$

- It has 5 variables, 3 pivots, and thus $5 - 3 = 2$ basic solutions
- To find a **particular solution**, remove the non-pivot columns, and (uniquely!) solve the resulting system:

$$\left(\begin{array}{ccc|c} \boxed{1} & 1 & 1 & 3 \\ 0 & \boxed{1} & 3 & 1 \\ 0 & 0 & \boxed{1} & 4 \end{array} \right)$$

- This has $(10, -11, 4)$ as solution; the original 5-variable system then has particular solution $(10, 0, -11, 0, 4)$.

Elaborated example, part II

- Consider the **associated homogeneous** system of equations:

$$\begin{array}{ccccc} x_1 & x_2 & x_3 & x_4 & x_5 \\ \begin{pmatrix} \boxed{1} & 1 & 1 & 1 & 1 \\ 0 & 0 & \boxed{1} & 2 & 3 \\ 0 & 0 & 0 & 0 & \boxed{1} \end{pmatrix} \end{array}$$

- The two **basic solutions** are found by removing each of the two non-pivot columns separately, and finding solutions:

$$\begin{array}{cccc} x_1 & x_3 & x_4 & x_5 \\ \begin{pmatrix} \boxed{1} & 1 & 1 & 1 \\ 0 & \boxed{1} & 2 & 3 \\ 0 & 0 & 0 & \boxed{1} \end{pmatrix} \end{array} \text{ and } \begin{array}{cccc} x_1 & x_2 & x_3 & x_5 \\ \begin{pmatrix} \boxed{1} & 1 & 1 & 1 \\ 0 & 0 & \boxed{1} & 3 \\ 0 & 0 & 0 & \boxed{1} \end{pmatrix} \end{array}$$

- We find: $(1, -2, 1, 0)$ and $(-1, 1, 0, 0)$. Adding zeros for missing columns gives: $(1, 0, -2, 1, 0)$ and $(-1, 1, 0, 0, 0)$.

Elaborated example, part III

Wrapping up: **all solutions** of the system

$$\left(\begin{array}{ccccc|c} \boxed{1} & 1 & 1 & 1 & 1 & 3 \\ 0 & 0 & \boxed{1} & 2 & 3 & 1 \\ 0 & 0 & 0 & 0 & \boxed{1} & 4 \end{array} \right)$$

are of the form:

$$\underbrace{(10, 0, -11, 0, 4)}_{\text{particular sol.}} + \underbrace{c_1(1, 0, -2, 1, 0) + c_2(-1, 1, 0, 0, 0)}_{\text{two basic solutions}}.$$

This is the **general solution** of the non-homogeneous system.

What are numbers?

Suppose I don't know what numbers are...
...but I still manage to pass Wiskundige Structuren.



Tell me: what are numbers?

What is the *first thing* you would tell me about some numbers, e.g. the real numbers?

What are numbers?

The First Thing: numbers form a **set**

S (← these are some numbers!)

The Second Thing: numbers can be **added together**

$$a \in S, b \in S \quad \Rightarrow \quad a + b \in S$$



Addition? Tell me more!


We have a set S , with a special operation '+' which satisfies:

1. $a + b = b + a$
2. $(a + b) + c = a + (b + c)$

...and there's a special element $\mathbf{0} \in S$ where:

3. $a + \mathbf{0} = a$

In math-speak, $(S, +, \mathbf{0})$ is called a **commutative monoid**, but we could also just call it a **set with addition**.



Examples: sets with addition

- Every kind of number you know: $\mathbb{R}, \mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{C}, \dots$
- The set of all polynomials:

$$(x^2 + 4x + 1) + (2x^2) := 3x^2 + 4x + 1 \quad \mathbf{0} := 0$$

- The set of all finite sets:

$$\{1, 2, 3\} + \{3, 4\} := \{1, 2, 3\} \cup \{3, 4\} = \{1, 2, 3, 4\} \quad \mathbf{0} := \{\}$$

- Here's a small example: $\{0\}$

$$0 + 0 := 0 \quad \mathbf{0} := 0$$

- ...and (important!) the set \mathbb{R}^n of all vectors of size n :

$$(x_1, \dots, x_n) + (y_1, \dots, y_n) := (x_1 + y_1, \dots, x_n + y_n) \quad \mathbf{0} := (0, \dots, 0)$$

Linear combinations

- We've been talking a lot about **linear combinations**:

$$a \cdot \mathbf{v} + b \cdot \mathbf{w} = \mathbf{u}$$

- **Q:** what is the most general kind of **set**, where we can take **linear combinations** of elements?
- **A:** a set V with addition and...**scalar multiplication**

$$a \in \mathbb{R}, \mathbf{v} \in V \quad \implies \quad a \cdot \mathbf{v} \in V$$



Multiplication?! What does that do?

A **vector space** is a set with addition $(V, +, \mathbf{0})$ with an extra operation \cdot , which satisfies:

- 1 $a \cdot (\mathbf{v} + \mathbf{w}) = a \cdot \mathbf{v} + a \cdot \mathbf{w}$
- 2 $(a + b) \cdot \mathbf{v} = a \cdot \mathbf{v} + b \cdot \mathbf{v}$
- 3 $a \cdot (b \cdot \mathbf{v}) = ab \cdot \mathbf{v}$
- 4 $1 \cdot \mathbf{v} = \mathbf{v}$
- 5 $0 \cdot \mathbf{v} = \mathbf{0}$

Example

Our **main example** is \mathbb{R}^n , where:

$$a \cdot (v_1, \dots, v_n) := (av_1, \dots, av_n)$$

Vector spaces: all together

Definition

A **vector space** $(V, +, \cdot, \mathbf{0})$ is a set V with a special element $\mathbf{0} \in V$ and operations '+' and ' \cdot ' satisfying:

① $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$

② $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$

③ $\mathbf{v} + \mathbf{0} = \mathbf{v}$

④ $a \cdot (\mathbf{v} + \mathbf{w}) = a \cdot \mathbf{v} + a \cdot \mathbf{w}$

⑤ $(a + b) \cdot \mathbf{v} = a \cdot \mathbf{v} + b \cdot \mathbf{v}$

⑥ $a \cdot (b \cdot \mathbf{v}) = ab \cdot \mathbf{v}$

⑦ $1 \cdot \mathbf{v} = \mathbf{v}$

⑧ $0 \cdot \mathbf{v} = \mathbf{0}$

for all $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$ and $a, b \in \mathbb{R}$.

Vector spaces: Main Example

Our **main example**:

$$\begin{aligned}\mathbb{R}^n &= \{(v_1, \dots, v_n) \mid v_1, \dots, v_n \in \mathbb{R}\} \\ &= \left\{ \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} \mid v_1, \dots, v_n \in \mathbb{R} \right\}\end{aligned}$$



The operations:

$$\begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} + \begin{pmatrix} w_1 \\ \vdots \\ w_n \end{pmatrix} = \begin{pmatrix} v_1 + w_1 \\ \vdots \\ v_n + w_n \end{pmatrix} \quad a \cdot \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} = \begin{pmatrix} av_1 \\ \vdots \\ av_n \end{pmatrix}$$




have a clear geometric interpretation.

Vector spaces: geometric interpretation

$a \cdot \mathbf{v}$ makes a vector shorter or longer:



$\mathbf{v} + \mathbf{w}$ stacks vectors together:



Example: subspaces

Certain **subsets** $V \subseteq \mathbb{R}^n$ are also vector spaces, e.g.

$$V = \{(v_1, v_2, 0) \mid v_1, v_2 \in \mathbb{R}\} \subseteq \mathbb{R}^3$$

$$W = \{(x, 2x) \mid x \in \mathbb{R}\} \subseteq \mathbb{R}^2$$

as long as they have $\mathbf{0}$, and they are **closed** under '+' and '·':

$$\mathbf{v}, \mathbf{w} \in V \implies \mathbf{v} + \mathbf{w} \in V$$

$$\mathbf{v} \in V, a \in \mathbb{R} \implies a \cdot \mathbf{v} \in V$$

These are called *subspaces* of \mathbb{R}^n .



Vector space example

We've seen this example before!

Example

The set of solutions of a homogeneous system of equations is a vector space.

Let S be the set of solutions of a homogeneous system of equations, with n variables. Then $S \subseteq \mathbb{R}^n$, and as we learned last week:

$$\mathbf{s}, \mathbf{t} \in S \implies \mathbf{s} + \mathbf{t} \in S$$

$$\mathbf{s} \in S, a \in \mathbb{R} \implies a \cdot \mathbf{s} \in S$$



Vector spaces: 'weirder' examples

\mathbb{R}^n and $V \subseteq \mathbb{R}^n$ are the only things we'll use in this course...but there are other examples:

- $\{0\}$ is still an example
- Polynomials are still an example: $5 \cdot (2x^2 + 1) = 10x^2 + 5$
- ...but finite sets are not!

$$5 \cdot \{\text{sandwich, Tuesday}\} = ???$$

- Functions $\mathcal{F}(X) := \{f : X \rightarrow \mathbb{R}\}$ are an example. If f, g are functions, then ' $f + g$ ' and $a \cdot f$ are also functions, defined by:

$$(f + g)(x) := f(x) + g(x) \qquad (a \cdot f)(x) = af(x)$$

Exercise: show that, if $X = \{1, 2, \dots, n\}$, then $\mathcal{F}(X)$ is basically the same as \mathbb{R}^n .



Our first theorem about vector spaces

We've got a **Definition**, let's prove a **Theorem**!

Theorem

Vector spaces have additive inverses. That is, for all $\mathbf{v} \in V$, there exists a vector $-\mathbf{v}$ such that $-\mathbf{v} + \mathbf{v} = \mathbf{0}$.

Proof. Let $-\mathbf{v} := (-1) \cdot \mathbf{v}$. Then, we use rules (1)-(6):

$$\begin{aligned} -\mathbf{v} + \mathbf{v} &= (-1) \cdot \mathbf{v} + 1 \cdot \mathbf{v} \\ &= (-1 + 1) \cdot \mathbf{v} \\ &= 0 \cdot \mathbf{v} \\ &= \mathbf{0} \end{aligned}$$